arXiv:0708.3114v4 [math.KT] 28 Sep 2008

DIFFERENTIAL TWISTED K-THEORY AND APPLICATIONS

ALAN L. CAREY, JOUKO MICKELSSON, AND BAI-LING WANG

Abstract. In this paper, we develop differential twisted K-theory and define a twisted Chern character on twisted K-theory which depends on a choice of connection and curving on the twisting gerbe. We also establish the general Riemann-Roch theorem in twisted K-theory and find some applications in the study of twisted K-theory of compact simple Lie groups.

Contents

1. Introduction 1
2. Preliminary notions 4
2.1. Twisted K-theory 4
2.2. Geometry of bundle gerbes 6
3. Differential twisted K-theory 8
3.1. The geometric model of differential twisted K-theory 9
3.2. Classifying space of twisted K-theory 17
4. Riemann-Roch theorem in twisted K-theory 22
5. Applications 26
5.1. The $SU(2)$ case 27
5.2. The $SU(3)$ case 30
References 32

1. Introduction

Generalized differential cohomology theories have recently excited considerable interest. For example, Cheeger-Simons differential characters play a role in index theory. In [27], Lott developed $\mathbb{R}/\mathbb{Z}$-valued index theory for Dirac operators coupled to virtual complex vector bundles with trivial Chern character, extending Atiyah-Patodi-Singer's reduced eta-invariants for flat vector bundles. The resulting K-theory is called K-theory with $\mathbb{R}/\mathbb{Z}$ coefficients. It is related to what is now called 'differential K-theory' as proposed by Freed [20], Hopkins and Singer [24], and further developed in [12]. Our aim here is to extend some of these ideas to create twisted differential K-theory.

Henceforth X will always denote a smooth manifold, $\mathcal{H}$ an infinite dimensional separable complex Hilbert space and $\mathbf{PU}(\mathcal{H})$ the projective unitary group of $\mathcal{H}$. The term 'twist' in

---

2000 *Mathematics Subject Classification.* 55R65, 53C29, 57R20, 81T13.

Key words and phrases. Differential twisted K-theory, Chern character, D-brane charges.

twisted K-theory $K^{ev/odd}(X,\sigma)$ of X refers to a continuous map $\sigma : X \to K(\mathbb{Z},3)$ which necessarily determines a principal $\mathbf{PU}(\mathcal{H})$-bundle $\mathcal{P}_\sigma$ over X.

We have been motivated to write this paper by some questions from physics. The reader unfamiliar with the language can pass on to the next paragraph. K-theory has previously been used to study anomaly cancellation problems for action principles in the presence of D-branes in string theory and M-theory. For Type II superstring theory with non-trivial B-field on X, it is believed that Ramond-Ramond charges lie in twisted K-theory $K^{ev/odd}(X,\sigma)$ of X (Cf.[39] [40]) with the twist, as above, given by $\sigma : X \to K(\mathbb{Z},3)$. The cohomology classes of Ramond-Ramond charges are twisted cohomology classes.

To describe this twisted cohomology we start with a closed differential form H representing the image of $[\sigma]$ under the map $H^3(X,\mathbb{Z}) \to H^3(X,\mathbb{R})$. We then need some additional concepts, reviewed in Section 2, beginning with the canonical 'lifting bundle gerbe' $\mathcal{G}_\sigma$ over X determined by $\mathcal{P}_\sigma$. It may be equipped with a gerbe connection θ and curving ω. Then H is in fact the normalized curvature of the triple $\check{\sigma} = (\mathcal{G}_\sigma, \theta, \omega)$. Next, twisted cohomology $H^{ev/odd}(X, d-H)$ is the cohomology of the complex of differential forms on X with the coboundary operator given by $d-H$. It is known (Cf. [4], [7], [28]) that the twisted Chern character

$$Ch_{\check{\sigma}} : K^{ev/odd}(X,\sigma) \longrightarrow H^{ev/odd}(X, d-H).$$

depends on a choice of a gerbe connection and a curving on the underlying gerbe $\mathcal{G}_\sigma$.

In many applications, the twisted Chern character is trivial due to the fact that the twisted K-groups are torsion. This means that the corresponding twisted cohomology $H^*(X, d-H)$ is zero in these examples. In order to study torsion elements in twisted K-theory, we give a geometric construction of a 'differential twisted K-theory with twisting given by $\check{\sigma}$'. That is, we replace ordinary twisted K-theory by a more refined cohomology theory that depends not only on the twist σ but also on the gerbe connection and curving. More specifically, in Section 3, we will construct our differential twisted K-theory, denoted $\check{K}^{ev/odd}(X,\check{\sigma})$ and a differential twisted Chern character $ch_{\check{\sigma}}$ on $\check{K}^{ev/odd}(X,\check{\sigma})$. We show that it gives a well-defined map

$$Ch_{\check{\sigma}} : K^{ev/odd}(X,\sigma) \longrightarrow H^{ev/odd}(X, d-H), \tag{1.1}$$

such that the following diagram commutes:

$$\begin{tikzcd}
\check{K}^{ev/odd}(X,\check{\sigma}) \arrow[r] \arrow[d, "ch_{\check{\sigma}}"] & K^{ev/odd}(X,\sigma) \arrow[d, "Ch_{\check{\sigma}}"] \\
\Omega_0^{ev/odd}(X, d-H) \arrow[r] & H^{ev/odd}(X, d-H),
\end{tikzcd} \tag{1.2}$$

where we have used the notation $\Omega_0^{ev/odd}(X, d-H)$ to denote the image of the differential Chern character map

$$ch_{\check{\sigma}} : \check{K}^{ev/odd}(X,\check{\sigma}) \longrightarrow \Omega^{ev/odd}(X, d-H).$$

In fact, the commutative diagram (1.2) is part of a commutative interlocking diagram of exact sequences which characterizes differential twisted K-theory $\check{K}^{ev/odd}(X, \check{\sigma})$, see Theorems 3.6 and 3.7 and Remark 3.8.

We remark that in the construction we need to choose local trivializations of the principal $\mathbf{PU}(\mathcal{H})$-bundle $\mathcal{P}_\sigma$. In addition we also need the notion of a 'spectral cut' (defined later on) for a $\mathbf{PU}(\mathcal{H})$-equivariant family of self-adjoint operators. Proposition 3.3 ensures that differential twisted K-theory and its differential Chern character don't depend on these choices. We also give a universal model of differential twisted K-theory (cf. Theorems 3.9 and 3.12) such that differential Chern character can be defined directly by a choice of a connection on a certain universal bundle classifying twisted K-theory.

When the twisting $\sigma : X \to K(\mathbb{Z}, 3)$ is trivial our construction gives rise to a different model for the ordinary differential K-theory previously studied in [12], [20] and [24]. Roughly stated the main idea is to provide some 'geometric cycles' as models for differential K-theory. Hopkins-Singer ([20], [24]) apply homotopy theory to the space of differential functions on the products of X with a varying simplex, taking values in a classifying space for K-theory. The viewpoint of [12] is to apply the family index theorem, so their models are geometric fibrations equipped with elliptic differential operators. We have utilized the spectral theory of self-adjoint operators to get local bundles over a groupoid, the double intersections of an open cover, for the odd case of twisted differential K-theory (and an adaptation of this for the even case).

In Section 4 we establish a generalization of the Atiyah-Hirzebruch Riemann-Roch theorem in twisted K-theory. Note that the Riemann-Roch theorem in twisted K-theory for K-oriented maps was discussed in [9]. Our refinement in Section 4 is to extend the theorem (see Theorems 4.1 and Theorem 4.4) so that it applies to non-K-oriented smooth maps $f : X \to Y$. When $[\sigma] = 0$ and f is K-oriented, our theorem reduces to the standard Atiyah-Hirzebruch Riemann-Roch theorem ([2]) .

In order to apply these ideas to string geometry we need one further concept, namely, the notion of twisted eta forms for those twisted K-classes that are torsion. This is useful in studying non-equivariant twisted K-theory of a compact Lie group. We discover that these twisted eta forms can be used to distinguish different twisted K-classes. Explicit computations are done for the Lie groups $SU(2)$ and $SU(3)$ as these examples arise in conformal field theory.

Acknowledgements. We thank the referee for carefully reading the manuscript and providing helpful advice. We also thank Thomas Schick for his comments on the manuscript. We acknowledge the assistance of Michael Murray for the discussion of Deligne cohomology. The authors acknowledge the support of: the Erwin Schrödinger Institute (AC, JM, BW), the Australian Research Council (AC, BW), the Clay Mathematics Institute (AC) and the Academy of Finland grant 516 8/07-08 (JM).

2. Preliminary notions

2.1. **Twisted K-theory.** In this subsection, we briefly review some basic facts about twisted K-theory, the main references are [3], [7], [16] and [34]. Let $\mathbf{PU}(\mathcal{H})$ denote the projective unitary group (equipped with the norm topology) of an infinite dimensional separable complex Hilbert space $\mathcal{H}$ and recall (cf. [25]) that it has the homotopy type of an Eilenberg-MacLane space $K(\mathbb{Z}, 2)$. The classifying space $B\mathbf{PU}(\mathcal{H})$ for principal $\mathbf{PU}(\mathcal{H})$-bundles is an Eilenberg-MacLane space $K(\mathbb{Z}, 3)$. Thus, the set of isomorphism classes of principal $\mathbf{PU}(\mathcal{H})$-bundles over X is canonically identified with the space $[X, K(\mathbb{Z}, 3)]$ of homotopy classes of maps from X to $K(\mathbb{Z}, 3)$ which in turn (Proposition 2.1 in [3]) is just $H^3(X, \mathbb{Z})$.

Much of the theory applies to locally compact, metrizable and separable spaces X, however, we will remain with the case where X is a manifold. Let $\mathcal{P}_\sigma$ be a principal $\mathbf{PU}(\mathcal{H})$-bundle over X whose classifying map $\sigma : X \to K(\mathbb{Z}, 3)$, called a twisting, defines an element $[\sigma] \in H^3(X, \mathbb{Z})$. There is a well-defined conjugation action of $\mathbf{PU}(\mathcal{H})$ on the space $\mathbf{Fred}(\mathcal{H})$ of bounded Fredholm operators. Let $\mathbf{Fred}(\mathcal{P}_\sigma) = \mathcal{P}_\sigma \times_{\mathbf{PU}(\mathcal{H})} \mathbf{Fred}(\mathcal{H})$ be the associated bundle of Fredholm operators.

Definition 2.1. If X is compact, the twisted K-group $K^0(X, \sigma)$ is defined to be the space of homotopy classes of sections of $\mathbf{Fred}(\mathcal{P}_\sigma)$ (see [34]). If X is locally compact, then the twisted K-group $K^0(X, \sigma)$ is defined to be the space of homotopy classes of 'compactly supported sections' of $\mathbf{Fred}(\mathcal{P}_\sigma)$, where compactly supported means these sections take values in the invertible operators away from a compact set. If X_0 is a closed subset of X, we may define in the obvious way the relative twisted K-group

$$K^0(X, X_0; \sigma) = K^0(X - X_0, \sigma).$$

Replacing $\mathbf{Fred}(\mathcal{H})$ by the space $\mathbf{Fred}^{sa}_*$ of self-adjoint Fredholm operators with both positive and negative essential spectrum, we can similarly define the twisted K-group $K^1(X, \sigma)$.

Remark 2.2. In the applications we consider unbounded self adjoint Fredholm operators of the form $D+A$ where D is a fixed unbounded self adjoint operator and A is bounded and varies. We utilise the topology on the unbounded self adjoint Fredholm operators induced by the map $D+A \to F_{D+A} = (D+A)(1+(D+A)^2)^{-1/2}$ [14]. This is justified for example by Theorem 8 in Appendix A of [14] which proves the estimate $||F_{D+A} - F_D|| < ||A||$. This estimate implies that if A varies smoothly in the uniform norm then so does F_{D+A} and hence in this sense so does $D + A$.

Next we choose a local trivialization of $\mathcal{P}_\sigma$ with respect to a good open cover $X = \bigcup_i U_i$ with transition functions given by $g_{ij} : U_i \cap U_j \longrightarrow \mathbf{PU}(\mathcal{H})$ and lifts $\hat{g}_{ij} : U_i \cap U_j \longrightarrow \mathbf{U}(\mathcal{H})$, for each g_{ij}. Then we have

$$\sigma_{ijk} \cdot Id = \hat{g}_{ij}\hat{g}_{jk}\hat{g}_{ki} \tag{2.1}$$

for a $U(1)$-valued Čech cocycle $\sigma_{ijk} : U_{ijk} \to U(1)$, where $U_{ijk} = U_i \cap U_j \cap U_k$ and Id is the identity operator in $\mathcal{H}$.

Then an element in $K^0(X,\sigma)$ can be represented by either a continuous $\mathbf{PU}(\mathcal{H})$-equivariant map $f:\mathcal{P}_\sigma\to\mathbf{Fred}(\mathcal{H})$, or by a twisted family of locally defined functions $\{f_i:U_i\longrightarrow\mathbf{Fred}(\mathcal{H})\}$ satisfying $f_i=\hat{g}_{ij}^{-1}f_j\hat{g}_{ij}$. Similarly, an element in $K^1(X,\sigma)$ can be represented either by a continuous $\mathbf{PU}(\mathcal{H})$-equivariant map $f:\mathcal{P}_\sigma\to\mathbf{Fred}^{sa}_*$, or a twisted family of locally defined functions $\{f_i:U_i\longrightarrow\mathbf{Fred}^{sa}_*\}$, satisfying $f_j=\hat{g}_{ij}^{-1}f_i\hat{g}_{ij}$.

Let $\sigma_1,\sigma_2:X\to K(\mathbb{Z},3)$ be a pair of continuous maps. We need to define a map $\sigma_1+\sigma_2$ from X to $K(\mathbb{Z},3)$ such that

$$[\sigma_1+\sigma_2]=[\sigma_1]+[\sigma_2]$$

in $H^3(X,\mathbb{Z})$. In order to define this map, we fix an isomorphism $\mathcal{H}\otimes\mathcal{H}\cong\mathcal{H}$ which induces a group homomorphism $\mathbf{U}(\mathcal{H})\times\mathbf{U}(\mathcal{H})\longrightarrow\mathbf{U}(\mathcal{H})$ whose restriction to the center is the group multiplication on $U(1)$. So we have a group homomorphism

$$\mathbf{PU}(\mathcal{H})\times\mathbf{PU}(\mathcal{H})\longrightarrow\mathbf{PU}(\mathcal{H})$$

which defines a continuous map, denoted m_*, of CW-complexes

$$B\mathbf{PU}(\mathcal{H})\times B\mathbf{PU}(\mathcal{H})\longrightarrow B\mathbf{PU}(\mathcal{H}).$$

As $B\mathbf{PU}(\mathcal{H})$ is a $K(\mathbb{Z},3)$, we may think of this as a continuous map taking $K(\mathbb{Z},3)\times K(\mathbb{Z},3)$ to $K(\mathbb{Z},3)$, which can be used to define

$$\sigma_1+\sigma_2:\quad X\xrightarrow{(\sigma_1,\sigma_2)}K(\mathbb{Z},3)\times K(\mathbb{Z},3)\xrightarrow{m_*}K(\mathbb{Z},3).$$

Just as in ordinary K-theory, we can define twisted K-groups $K^i(X,\sigma)$ for all $i\in\mathbb{Z}$ such that the twisted K-theory is a generalized cohomology theory with period 2 on the category of topological spaces equipped with a principal $\mathbf{PU}(\mathcal{H})$-bundle. Twisted K-theory satisfies the following basic properties [3][16].

(1) For any proper continuous map $f:X\to Y$ there exists a natural pull-back map

$$f^*:K^i(Y,\sigma)\longrightarrow K^i(X,f^*\sigma), \tag{2.2}$$

for any $\sigma:Y\to K(\mathbb{Z},3)$.

(2) With a fixed isomorphism $\mathcal{H}\otimes\mathcal{H}\to\mathcal{H}$, there is a homomorphism

$$K^i(X,\sigma_1)\times K^j(X,\sigma_2)\longrightarrow K^{i+j}(X,\sigma_1+\sigma_2)$$

for any two twistings $\sigma_1,\sigma_2:X\to K(\mathbb{Z},3)$.

(3) If X is covered by two closed subsets X_1 and X_2, there is a Mayer-Vietoris exact sequence

$$\begin{tikzcd}
K^1(X_1,\sigma_1)\oplus K^1(X_2,\sigma_2) \arrow[r] & K^1(X_1\cap X_2,\sigma_{12}) \arrow[r] & K^0(X,\sigma) \arrow[d] \\
K^1(X,\sigma) \arrow[u] & K^0(X_1\cap X_2,\sigma_{12}) \arrow[l] & K^0(X_1,\sigma_1)\oplus K^0(X_2,\sigma_2) \arrow[l]
\end{tikzcd}$$

where σ_1, σ_2 and σ_{12} are the restrictions of $\sigma:X\to K(\mathbb{Z},3)$ to X_1, X_2 and $X_1\cap X_2$ respectively.

(4) If X is covered by two open subsets U_1 and U_2, there is a Mayer-Vietoris exact sequence

$$\begin{tikzcd}
K^0(X,\sigma) \arrow[r] & K^1(U_1\cap U_2,\sigma_{12}) \arrow[r] & K^1(U_1,\sigma_1)\oplus K^1(U_2,\sigma_2) \arrow[d] \\
K^0(U_1,\sigma_1)\oplus K^0(U_2,\sigma_2) \arrow[u] & K^0(U_1\cap U_2,\sigma_{12}) \arrow[l] & K^1(X,\sigma) \arrow[l]
\end{tikzcd}$$

where σ_1, σ_2 and σ_{12} are the restrictions of $\sigma : X \to K(\mathbb{Z},3)$ to U_1, U_2 and $U_1\cap U_2$ respectively.

(5) (**Thom isomorphism**) Let $\pi : E \to X$ be an oriented real vector bundle of rank k over X with the classifying map denoted by $\nu_E : X \to \mathbf{BSO}(k)$, then there is a canonical isomorphism, for any twisting $\alpha : X \to K(\mathbb{Z},3)$,

$$K^{ev/odd}(X,\alpha + W_3\circ\nu_E) \cong K^{ev/odd}(E,\alpha\circ\pi), \tag{2.3}$$

with the grading shifted by $k(mod\ 2)$. Here $W_3 : \mathbf{BSO}(k) \to K(\mathbb{Z},3)$ is the classifying map of the principal $K(\mathbb{Z},2)$-bundle $\mathbf{BSpin}^c(k) \to \mathbf{BSO}(k)$.

(6) For any differentiable map $f : X \to Y$ between two smooth manifolds X and Y, there is a natural push-forward map

$$f_!^K : K^i\big(X, f^*\sigma + W_3\circ\nu_f\big) \longrightarrow K^{i+d(f)}(Y,\sigma), \tag{2.4}$$

for any $\sigma \in H^3(Y,\mathbb{Z})$, $d(f) = \dim(X) - \dim(Y) \mod 2$, and $\nu_f : X \to \mathbf{BSO}$ is a classifying map of $TX \oplus f^*TY$. Note that $W_3\circ\nu_f$ is a map $X \to K(\mathbb{Z},3)$ representing the image of $w_2(X) + f^*(w_2(Y)) \in H^2(X,\mathbb{Z}_2)$ under the Bockstein homomorphism $H^2(X,\mathbb{Z}_2) \longrightarrow H^3(X,\mathbb{Z})$.

2.2. **Geometry of bundle gerbes.** In this subsection, we recall some basics of bundle gerbes and their connections and curvings from [31, 32]. First recall that associated to the central extension

$$1 \to U(1) \longrightarrow U(\mathcal{H}) \longrightarrow \mathbf{PU}(\mathcal{H}) \to 1. \tag{2.5}$$

there exists a so-called lifting bundle gerbe $\mathcal{G}_\sigma$ [31]. This object is represented diagrammatically by

$$\begin{tikzcd}
\mathcal{G}_\sigma \arrow[d] & & . \\
\mathcal{P}_\sigma^{[2]} \arrow[r, shift left, "\pi_1"] \arrow[r, shift right, "\pi_2"'] & \mathcal{P}_\sigma \arrow[d, "\pi"] & \\
 & X &
\end{tikzcd} \tag{2.6}$$

This diagram should be read as follows. Starting with $\pi : \mathcal{P}_\sigma \to X$ form the fibre product $\mathcal{P}_\sigma^{[2]}$ which is a groupoid $\mathcal{P}_\sigma^{[2]} = \mathcal{P}_\sigma \times_X \mathcal{P}_\sigma$ with source and range maps $\pi_1 : (y_1,y_2) \mapsto y_1$ and $\pi_2 : (y_1,y_2) \mapsto y_2$. There is an obvious map from each fiber of $\mathcal{P}_\sigma^{[2]}$ to $\mathbf{PU}(\mathcal{H})$ and so we can define the fiber of $\mathcal{G}_\sigma$ over a point in $\mathcal{P}_\sigma^{[2]}$ by pulling back the fibration (2.5) using this map. This endows $\mathcal{G}_\sigma$ with a groupoid structure (from the multiplication in $U(\mathcal{H})$) and in fact it is a $U(1)$-groupoid extension of $\mathcal{P}_\sigma^{[2]}$.

We will review the notion of a bundle gerbe connection, curving and curvature on $\mathcal{G}_\sigma$. Recall that there is an exact sequence of differential p-forms

$$\Omega^p(X) \xrightarrow{\pi^*} \Omega^p(\mathcal{P}_\sigma) \xrightarrow{\delta} \Omega^p(\mathcal{P}_\sigma^{[2]}) \xrightarrow{\delta} \cdots, \tag{2.7}$$

Here $\mathcal{P}_\sigma^{[q]}$ is the q-fold fiber product of π, $\delta = \sum_{i=1}^{q+1}(-1)^i \pi_i^* : \Omega^p(\mathcal{P}_\sigma^{[q]}) \to \Omega^p(\mathcal{P}_\sigma^{[q+1]})$ is the alternating sum of pull-backs of projections, where π_i is the projection map which omits the i-th point in the fibre product.

A gerbe connection on $\mathcal{P}_\sigma$ is a unitary connection θ on the principal $U(1)$-bundle $\mathcal{G}_\sigma$ over $\mathcal{P}_\sigma$ which commutes with the bundle gerbe product. A bundle gerbe connection θ has curvature

$$F_\theta \in \Omega^2(\mathcal{P}_\sigma)$$

satisfying $\delta(F_\theta) = 0$ and hence from the exact sequence (2.7) for $p = 2$, there exists a two-form ω on $\mathcal{P}_\sigma$ such that

$$F_\theta = \pi_2^*(\omega) - \pi_1^*(\omega).$$

Such an ω is called a curving for the gerbe connection θ. The choice of a curving is not unique, the ambiguity in the choice is precisely the addition of the pull-back to $\mathcal{P}_\sigma$ of a two-form on X. Given a choice of curving ω we have

$$\delta(d\omega) = d\delta(\omega) = dF_\theta = 0$$

so that we can find a unique closed three-form on β on X, such that $d\omega = \pi^*\beta$. We denote by $\check{\sigma} = (\mathcal{G}_\sigma, \theta, \omega)$ the lifting bundle gerbe $\mathcal{G}_\sigma$ with the connection θ and a curving ω. Moreover $H = \dfrac{\beta}{2\pi\sqrt{-1}}$ is a de Rham representative for the Dixmier-Douady class $[\sigma]$. We call H the normalized curvature of $\check{\sigma}$.

Consider a good open cover $\mathcal{U} = \{U_i\}$ of X such that $\mathcal{P}_\sigma \to X$ has trivializing sections ϕ_i over each U_i with transition functions $g_{ij} : U_i \cap U_j \longrightarrow \mathbf{PU}(\mathcal{H})$ satisfying $\phi_j = \phi_i g_{ij}$ over $U_i \cap U_j$. Define σ_{ijk} by $\hat{g}_{ij}\hat{g}_{jk} = \hat{g}_{ik}\sigma_{ijk}$ for a lift of g_{ij} to $\hat{g}_{ij} : U_i \cap U_j \to \mathbf{U}(\mathcal{H})$. Note that the pair (ϕ_i, ϕ_j) defines a section of $\mathcal{P}_\sigma^{[2]}$ over $U_i \cap U_j$. The connection θ can be pulled back by (ϕ_i, ϕ_j) to define a 1-form A_{ij} on $U_i \cap U_j$ and the curving ω can be pulled-back by the ϕ_i to define two-forms B_i on U_i. Then the triple

$$(\sigma_{ijk}, A_{ij}, B_i) \tag{2.8}$$

is a degree two smooth Deligne cocycle.

In our construction of the Chern character for differential twisted K-theory and in our proof of the Riemann-Roch theorem for twisted K-theory we need to work with the Deligne data defined by $\check{\sigma}$. So we recall for the reader's convenience that degree p smooth Deligne cohomology is the p-th Čech hypercohomology group of the complex of sheaves on X (Cf. [11]):

$$\underline{U(1)} \xrightarrow{d\log} \Omega^1_X \xrightarrow{d} \Omega^2_X \longrightarrow \cdots \xrightarrow{d} \Omega^p_X$$

where $\underline{U(1)}$ is the sheaf of germs of smooth $U(1)$-valued functions on M and Ω^p_X is the sheaf of germs of imaginary-valued differential p-forms on X. A degree 0 smooth Deligne class is represented by a smooth map $f\colon X \to U(1)$. A degree 1 smooth Deligne class ξ can be represented by Hermitian line bundles with Hermitian connection.

The degree 2 Deligne cohomology group $H^2_{\mathcal{D}}(X)$ can be calculated as the cohomology of the total complex of the double complex with respect to a good cover $\mathcal{U} = \{U_i\}$ of X:

(2.9)
$$\begin{tikzcd}
\vdots & \vdots & \vdots \\
C^2(\mathcal{U}, \underline{U(1)}) \arrow[r, "d\log"] \arrow[u, "\delta"] & C^2(\mathcal{U}, \Omega^1_X) \arrow[r, "d"] \arrow[u, "\delta"] & C^2(\mathcal{U}, \Omega^2_X) \arrow[u, "\delta"] \\
C^1(\mathcal{U}, \underline{U(1)}) \arrow[r, "d\log"] \arrow[u, "\delta"] & C^1(\mathcal{U}, \Omega^1_X) \arrow[r, "d"] \arrow[u, "\delta"] & C^1(\mathcal{U}, \Omega^2_X) \arrow[u, "\delta"] \\
C^0(\mathcal{U}, \underline{U(1)}) \arrow[r, "d\log"] \arrow[u, "\delta"] & C^0(\mathcal{U}, \Omega^1_X) \arrow[r, "d"] \arrow[u, "\delta"] & C^0(\mathcal{U}, \Omega^2_X) \arrow[u, "\delta"]
\end{tikzcd}$$

which is the quotient of the abelian group of degree two Deligne cocycles by the subgroup of degree two Deligne coboundaries. Here a triple

$$(\sigma_{ijk}, A_{ij}, B_i) \in C^2(\mathcal{U}, \underline{U(1)}) \oplus C^1(\mathcal{U}, \Omega^1_X) \oplus C^0(\mathcal{U}, \Omega^2_X)$$

is a Deligne cocycle if it satisfies the following cocycle condition

1. $\sigma_{ijk}\sigma^{-1}_{ijl}\sigma_{ikl}\sigma^{-1}_{jkl} = 1$.
2. $A_{ij} + A_{jk} + A_{ki} = d\log(\sigma_{ijk})$.
3. $B_j - B_i = dA_{ij}$.

A degree two Deligne coboundary is a triple of the following type

$$(h_{ij}h^{-1}_{ik}h_{jk}, d\log(h_{ij}) + a_i - a_j, da_i)$$

for $(h_{ij}, a_i) \in C^1(\mathcal{U}, \underline{U(1)}) \oplus C^0(\mathcal{U}, \Omega^1_X)$. The degree 2 Deligne cohomology of X classifies stable isomorphism classes of Hermitian bundle gerbes with connection and curving (see [32]).

We conclude this subsection by noting an important consequence of the above discussion namely that the degree two Deligne class of (2.8) determined by $\check{\sigma} = \{\mathcal{G}_\sigma, \theta, \omega)$ is independent of the choice of locally trivializing sections of $\mathcal{P}_\sigma$ and the lifting $\hat{g}_{ij}$.

3. Differential twisted K-theory

Recall our data: X is a compact smooth manifold, $\mathcal{P}_\sigma$ is a principal $\mathbf{PU}(\mathcal{H})$ bundle over X whose classifying map is given by $\sigma : X \longrightarrow K(\mathbb{Z}, 3)$. There is a lifting bundle gerbe $\mathcal{G}_\sigma$ with a bundle gerbe connection θ and a curving ω with $\check{\sigma} = (\mathcal{G}_\sigma, \theta, \omega)$ denoting this triple. We refer to $\check{\sigma}$ as a twisting in differential twisted K-theory. The normalized curvature of the bundle gerbe with connection and curving $\check{\sigma}$ is denoted by H. With respect to an open cover $\{U_i\}$ of X which trivializes $\mathcal{P}_\sigma$ using local sections $\{\phi_i\}$, we have an associated Deligne 2-cocycle $(\sigma_{ijk}, A_{ij}, B_i)$.

We will establish in the next subsection the existence of a differential twisted K-theory and construct its differential twisted Chern character which induces the twisted Chern character $ch_{\check{\sigma}} : K^*(X, \check{\sigma}) \longrightarrow H^*(X, d - H)$.

3.1. **The geometric model of differential twisted K-theory.** We begin with odd twisted K-theory. Let $f : \mathcal{P}_\sigma \to \mathbf{Fred}_*^{sa}$ be a $\mathbf{PU}(\mathcal{H})$ equivariant map to the bounded self-adjoint Fredholm operators with both positive and negative essential spectrum, i.e., the homotopy class of f is an element of $K^1(X, \sigma)$. We make the additional assumption that the operators $f(x)$ have discrete spectrum (which is the case in many physics examples). This is no real limitation, since by [5] the space $\mathbf{Fred}_*^{sa}$ is homotopy equivalent to the subspace $\mathcal{F}_*^{sa}$ of operators of norm less than or equal to one and with essential spectrum ± 1. Moreover the map from unbounded to bounded Fredholms $D \to D(1 + D^2)^{-1/2}$ introduced earlier maps the unbounded self adjoint operators with discrete spectrum into the space $\mathcal{F}_*^{sa}$.

Next choose an open cover $\{U_i\}$, $i = 1, 2, \ldots, n$, of X such that on each U_i there is a local section $\phi_i : U_i \to \mathcal{P}_\sigma$ and for each i there is a real number λ_i not in the spectrum of the operators $f(\phi_i(x))$, for all $x \in U_i$. We refer to λ_i as a choice of *spectral cut* for the family $\{f(\phi_i(x))\}_{x \in U_i}$. Furthermore, we can require that in each interval (λ_i, λ_j) there are only finite number of eigenvalues and each with finite multiplicity using the model $\mathcal{F}_*^{sa}$ above and selecting the λ_i's in the open interval $(-1, 1)$. Then over each $U_{ij} = U_i \cap U_j$ we have a finite rank vector bundle E_{ij} spanned by the eigenvectors of $f(\phi_j(x))$ with eigenvalues in the open interval (λ_i, λ_j), with a chosen ordering $\lambda_i < \lambda_j$ for $i < j$.

Let the transition functions be denoted by $g_{ij} : U_{ij} \to \mathbf{PU}(\mathcal{H})$ such that $\phi_j(x) = \phi_i(x) g_{ij}(x)$. If $\hat{g}_{ij}$ is a lift of g_{ij} to $\mathbf{U}(\mathcal{H})$ then $\hat{g}_{ij}\hat{g}_{jk}\hat{g}_{ki} = \sigma_{ijk} Id$ on triple intersections $U_{ijk} = U_i \cap U_j \cap U_k$ with σ_{ijk} taking values in $U(1)$.

Next we define bundle maps $\phi_{ijk} : E_{ij} \to E_{ik}$ over U_{ijk} for $\lambda_i < \lambda_j < \lambda_k$ as follows. Note that $f(\phi_j(x)) = \hat{g}_{kj}^{-1} f(\phi_k(x)) \hat{g}_{kj}$. First act by $\hat{g}_{kj}$ on E_{ij} and then use the inclusion to E_{ik}. The vector bundle $\phi_{ijk}(E_{ij})$ can be identified as the tensor product $L_{kj} \otimes E_{ij}$ where the complex line bundle L_{kj} over U_{kj} comes from the lifting bundle gerbe, i.e., from the pull-back of the central extension of $\mathbf{PU}(\mathcal{H})$ with respect to the map $g_{kj} : U_{jk} \to \mathbf{PU}(\mathcal{H})$. Thus we have

$$L_{kj} \otimes E_{ij} \oplus E_{jk} = E_{ik}, \tag{3.1}$$

over U_{ijk}. We call this the twisted cocycle property of families of local vector bundles $\{E_{ij}\}$.

In the untwisted case we can identify E_{ji} as the virtual bundle $-E_{ij}$. In the twisted case we have to remember that the vector bundles are defined using the local sections attached to the second index. Therefore we identify E_{ji} with $-L_{ji} \otimes E_{ij}$ the twist coming again from the transition function g_{ji} relating the local sections on open sets U_i, U_j.

Since the vector bundles E_{ij} are defined via projections P_{ij} onto finite dimensional subspaces $P_{ij}\mathcal{H}$, they come equipped with a natural connection ∇_{ij} and we can extend the above equality (3.1) to

$$(L_{kj}, A_{kj}) \otimes (E_{ij}, \nabla_{ij}) \oplus (E_{jk}, \nabla_{jk}) = (E_{ik}, \nabla_{ik}) \tag{3.2}$$

where A_{ij} is the gerbe connection determined by the differential twisting $\check{\sigma} = (\mathcal{G}_\sigma, \theta, \omega)$. For simplicity, we normalize B_i such that the first Chern class c_{ij} of L_{ij} is represented by $B_j - B_i$.

Remark 3.1. In the case of a trivial $\mathbf{PU}(\mathcal{H})$ bundle one has a choice of lifts such that $\sigma_{ijk} = 1$ and actually one has a global family of Fredholm operators parametrized by points on U_{ij}. In the untwisted case, the spectral subspaces E_{ij} are directly parametrized by points in X and we have $E_{ij} \oplus E_{jk} = E_{ik}$ over U_{ijk}.

Remark 3.2. There is an inverse map from the twisted cocycle of local vector bundles to a global object in $K^1(X, \sigma)$. To make the construction simple, we use the alternative classifying space $\mathbf{U}_1(\mathcal{H})$ of unitary operators which differ from the unit by a trace-class operator. Any vector bundle E_{ij} can be defined using a projection valued map $P_{ij} : U_{ij} \to L(\mathcal{H})$. We assume that the projections $P_{ij}(x)$ commute if the second indices are equal. (This is automatically the case when the projections are defined from a twisted K-theory element as above.) On the overlap U_{ijk} we require

$$\hat{g}_{jk}^{-1} P_{ij} \hat{g}_{jk} + P_{jk} = P_{ik}, \tag{3.3}$$

this being the twisted cocycle property (3.1) of the vector bundles $\{E_{ij}\}$. We put

$$g_i(x) = e^{2\pi i \sum_j \rho_j P_{ji}((x)}$$

where $\sum \rho_i(x) = 1$ is a partition of unity subordinate to the open cover $\{U_i\}$. The function $g_i : U_i \longrightarrow \mathbf{U}_1(\mathcal{H})$, satisfies

$$\hat{g}_{ji}^{-1} (\log\, g_j) \hat{g}_{ji} = 2\pi i\, \hat{g}_{ji}^{-1} \sum_k \rho_k P_{kj} \hat{g}_{ji}, = \log(g_i) - 2\pi i P_{ji}$$

by (3.3), which implies

$$\hat{g}_{ji}^{-1} g_j \hat{g}_{ji} = g_i e^{-2\pi i P_{ji}} = g_i$$

on the overlap U_{ij}. Hence, $\{g_i\}$ defines an element in $K^1(X, \sigma)$.

Let ℓ_{ij} be the top exterior power of E_{ij} and n_{ij} the rank of E_{ij}. Then the collection of integers $\{n_{ij}\}$ is an integral Čech cocycle. Furthermore,

$$L_{kj}^{n_{ij}} \otimes \ell_{ij} \otimes \ell_{jk} = \ell_{ik}. \tag{3.4}$$

As noted in [30] in the case when n_{ij} is trivial, i.e., $n_{ij} = n_i - n_j$ for some locally constant integer valued functions n_i, one can define $\ell'_{ij} = \ell_{ij} \otimes L_{ij}^{n_j}$ and one has

$$\ell'_{ij} \otimes \ell'_{jk} = \ell'_{ik}.$$

This gerbe is however defined only modulo integer powers of L since one can always shift $n_i \mapsto n_i + n$ for a constant n.

In any case, we obtain from (3.2) the cocycle relation for the Chern character forms of the vector bundles involved,

$$e^{B_j - B_k}\omega_{ij} + \omega_{jk} = \omega_{ik}, \tag{3.5}$$

where $\omega_{ij} = ch(E_{ij}, \nabla_{ij})$ is the Chern character form of (E_{ij}, ∇_{ij}).

Denote by $\bar{\omega}_{ij}$ the even differential form $e^{B_j}\omega_{ij}$. We have then

$$\delta(\bar{\omega}_{ij}) = \bar{\omega}_{jk} - \bar{\omega}_{ik} + \bar{\omega}_{ij} = 0, \qquad (d-H)\bar{\omega}_{ij} = 0.$$

Applying the tic-tac-toe argument to the following twisted Čech-de Rham double complex,

(3.6)

$$\begin{tikzcd}
{} & {} & {} & \\
\Omega^{even}(\{U_{ijk}\}) \arrow[u, "\delta"] \arrow[r, "d-H"] & \Omega^{odd}(\{U_{ijk}\}) \arrow[u, "\delta"] \arrow[r, "d-H"] & \Omega^{even}(\{U_{ijk}\}) \arrow[u, "\delta"] \arrow[r, "d-H"] & {} \\
\Omega^{even}(\{U_{ij}\}) \arrow[u, "\delta"] \arrow[r, "d-H"] & \Omega^{odd}(\{U_{ij}\}) \arrow[u, "\delta"] \arrow[r, "d-H"] & \Omega^{even}(\{U_{ij}\}) \arrow[u, "\delta"] \arrow[r, "d-H"] & {} \\
\Omega^{even}(\{U_{i}\}) \arrow[u, "\delta"] \arrow[r, "d-H"] & \Omega^{odd}(\{U_{i}\}) \arrow[u, "\delta"] \arrow[r, "d-H"] & \Omega^{even}(\{U_{i}\}) \arrow[u, "\delta"] \arrow[r, "d-H"] & {} \\
\Omega^{even}(X) \arrow[u, "\delta"] \arrow[r, "d-H"] & \Omega^{odd}(X) \arrow[u, "\delta"] \arrow[r, "d-H"] & \Omega^{even}(X) \arrow[u, "\delta"] \arrow[r, "d-H"] & {}
\end{tikzcd}$$

we obtain an odd-degree $(d-H)$-closed differential form Θ via the following diagram chase:

(3.7)

$$\begin{tikzcd}
0 & & \\
\bar{\omega}_{ij} \arrow[u, "\delta"] \arrow[r, "d-H"] & 0 & \\
\eta_i \arrow[u, "\delta"] \arrow[r] & (d-H)\eta_i \arrow[u, "\delta"] \arrow[r] & 0 \\
& \Theta \arrow[u, "\delta"] \arrow[r, "d-H"] & 0 \arrow[u, "\delta"]
\end{tikzcd}$$

with respect to an open cover $\{U_i\}$. Specifically, we can choose

$$\eta_i = \sum_k \rho_k \bar{\omega}_{ki}$$

satisfying $\delta(\eta_i) = \bar{\omega}_{ij}$ by direct calculation. Applying $d-H$ to η_i and diagram chasing, we have a locally defined odd differential form $(d-H)\eta_i$ on each U_i such that

$$(d-H)\eta_i = (d-H)\eta_j$$

over U_{ij}, hence we obtain a globally defined odd differential form Θ such that

$$\Theta|_{U_i} = (d-H)\eta_i, \qquad (d-H)\Theta = 0.$$

This globally defined odd differential form Θ is denoted by

$$ch_{\check\sigma}(f, \{\lambda_i\}, \{\rho_i\}),$$

and is called the twisted differential Chern character of f. Clearly it can depend on the choice of spectral cut $\{\lambda_i\}$ of $\{f\circ\phi_i\}$ associated to local trivializing sections $\{\phi_i : U_i \to \mathcal{P}_\sigma\}$, and on a choice of a partition of unity $\{\rho_i\}$ on X subordinate to $\{U_i\}$.

The next proposition explains how $ch_{\check\sigma}(f, \{\lambda_i\}, \{\rho_i\})$ changes if we change the choice of spectral cuts $\{\lambda_i\}$, or the partition of unity $\{\rho_i\}$ on X. We can also see the dependence on $\mathbf{PU}(\mathcal{H})$-equivariant homotopies.

Proposition 3.3. (1) *Choose local trivializing sections $\{\phi_i : U_i \to \mathcal{P}_\sigma\}$ and let $\{\lambda_i\}$ be a family of spectral cuts for $\{f\circ\phi_i\}$. Now let $\{\rho_i\}$ and $\{\rho'_i\}$ be a pair of partitions of unity on X subordinate to $\{U_i\}$, then there exists an even differential form $\eta(f, \{\lambda_i\}, \{\rho_i\}, \{\rho'_i\})$ on X such that*

$$ch_{\check\sigma}(f, \{\lambda_i\}, \{\rho_i\}) - ch_{\check\sigma}(f, \{\mu_i\}, \{\rho'_i\}) = (d-H)\eta(f, \{\lambda_i\}, \{\rho_i\}, \{\rho'_i\}).$$

(2) *Let $\{\lambda_i < \mu_i\}$ be a pair of spectral cuts for the family $\{f\circ\phi_i\}$ on an open cover $\{U_i\}$ of X. Then there exists an even differential form $\eta(f, \{\lambda_i\}, \{\mu_i\}, \{\rho_i\})$ on X such that*

$$ch_{\check\sigma}(f, \{\lambda_i\}, \{\rho_i\}) - ch_{\check\sigma}(f, \{\mu_i\}, \{\rho_i\}) = (d-H)\eta(f, \{\lambda_i\}, \{\mu_i\}, \{\rho_i\}).$$

(3) *Let $f : P_\sigma \times [t_0, t_1] \to \mathbf{Fred}^{sa}_*$ be a $\mathbf{PU}(\mathcal{H})$-equivariant homotopy connecting $f_0 = f(\cdot, t_0)$ and $f_1 = f(\cdot, t_1)$ such that there is a family of spectral cuts $\{\lambda_i\}$ independent of $t \in [t_0, t_1]$ for $\{f(\phi_i, t)\}$ on an open cover $\{U_i \times [t_0, t_1]\}$ of $X \times [t_0, t_1]$. Then there exists an even differential form $\eta(f_0, f_1, \{\lambda_i\}, \{\rho_i\})$ on X such that*

$$ch_{\check\sigma}(f_0, \{\lambda_i\}, \{\rho_i\}) - ch_{\check\sigma}(f_1, \{\lambda_i\}, \{\rho_i\}) = (d-H)\eta(f_0, f_1, \{\lambda_i\}, \{\rho_i\}).$$

Proof. (1) From the definition of the twisted differential Chern character, we have

$$ch_{\check\sigma}(f, \{\lambda_i\}, \{\rho_i\})|_{U_j} = \sum_i (d-H)\rho_i \wedge e^{B_j} ch(E_{ij}, \nabla_{ij})$$

for the spectral cut $\{\lambda_i\}$ and a partition of unity $\{\rho_i\}$, and

$$ch_{\check\sigma}(f, \{\lambda_i\}, \{\rho'_i\})|_{U_j} = \sum_i (d-H)\rho'_i \wedge e^{B_j} ch(E_{ij}, \nabla_{ij})$$

for the spectral cut $\{\lambda_i\}$ and a partition of unity $\{\rho'_i\}$. Their difference is given by

$$\sum_i (d-H)\big((\rho_i - \rho'_i)e^{B_j} ch(E_{ij}, \nabla_{ij})\big).$$

Note that $\sum_i \big((\rho_i - \rho'_i)e^{B_j} ch(E_{ij}, \nabla_{ij})\big)$ is a globally defined even differential form denoted by $\eta(f, \{\lambda_i\}, \{\rho_i\}, \{\rho'_i\})$, as

$$\sum_i \big((\rho_i - \rho'_i)e^{B_j} ch(E_{ij}, \nabla_{ij})\big) = \sum_i \big((\rho_i - \rho'_i)e^{B_k} ch(E_{ik}, \nabla_{ik})\big)$$

over $U_j \cap U_k$ by the cocycle condition of $\{e^{B_j} ch(E_{ij}, \nabla_{ij})\}$. This verifies that

$$ch_{\check\sigma}(f, \{\lambda_i\}, \{\rho_i\}) - ch_{\check\sigma}(f, \{\mu_i\}, \{\rho'_i\}) = (d-H)\eta(f, \{\lambda_i\}, \{\rho_i\}, \{\rho'_i\}).$$

(2) Let E_i be the vector bundle over U_i, equal to the spectral subspace defined by the open interval (λ_i, μ_i) in the spectrum of $f(\phi_i(x))$. Then we have

$$F_{ij} \oplus L_{ji} \otimes E_i = E_{ij} \oplus E_j$$

on U_{ij}. Equipped with the natural connections, we have

$$(F_{ij}, \nabla_{F_{ij}}) \oplus (L_{ij}, A_{ij}) \otimes (E_i, \nabla_{E_i}) \cong (E_{ij}, \nabla_{E_{ij}}) \oplus (E_j, \nabla_{E_j}).$$

Hence,

$$e^{B_j} ch(F_{ij}, \nabla_{F_{ij}}) = e^{B_j} ch(E_{ij}, \nabla_{E_{ij}}) + e^{B_j} ch(E_j, \nabla_{E_j}) - e^{B_i} ch(E_i, \nabla_{E_i}).$$

Note that

(a) $ch_{\check{\sigma}}(f, \{\lambda_i\}, \{\rho_i\})|_{U_j} = (d - H) \sum_i \rho_i e^{B_j} ch(E_{ij}, \nabla_{E_{ij}})$.
(b) $ch_{\check{\sigma}}(f, \{\mu_i\}, \{\rho_i\})|_{U_j} = (d - H) \sum_i \rho_i e^{B_j} ch(F_{ij}, \nabla_{F_{ij}})$.
(c) $e^{B_j} ch(E_j, \nabla_{E_j}) = \sum_i \rho_i e^{B_j} ch(E_j, \nabla_{E_j})$ is $d - H$ closed.

Then the difference of the twisted differential Chern character form of f with respect to a pair of spectral cuts $\{\lambda_i < \mu_i\}$ is given by

$$ch_{\check{\sigma}}(f, \{\lambda_i\}, \{\rho_i\}) - ch_{\check{\sigma}}(f, \{\mu_i\}, \{\rho_i\}) = (d - H)\eta(f, \{\lambda_i\}, \{\mu_i\}, \{\rho_i\})$$

where $\eta(f, \{\lambda_i\}, \{\mu_i\}, \{\rho_i\}) = \sum_i \rho_i e^{B_i} ch(E_i, \nabla_{E_i})$.

(3) Let $E_{ij}(t)$ be the finite rank vector bundle with the natural connection $\nabla(t)$ over $U_i \cap U_j$, whose fiber at x is spanned by the eigenvectors of $f(\phi_i(x), t)$ with eigenvalues in the open interval (λ_i, λ_j). Then $E_{ij}(t)$ is a continuous family of vector bundles over $U_i \cap U_j$ with natural connections satisfying

$$(L_{kj}, A_{kj}) \otimes (E_{ij}(t), \nabla_{ij}(t)) \oplus (E_{jk}(t), \nabla_{jk}(t)) = (E_{ik}(t), \nabla_{ik}(t)).$$

Applying the standard Chern-Weil argument to the odd differential form $ch_{\check{\sigma}}(f, \{\lambda_i\})$ on $M \times [t_0, t_1]$, we have

$$ch_{\check{\sigma}}(f_0, \{\lambda_i\}, \{\rho_i\}) - ch_{\check{\sigma}}(f_1, \{\lambda_i\}, \{\rho_i\}) = (d - H)\eta(f_0, f_1, \{\lambda_i\}, \{\rho_i\})$$

for $\eta(f_0, f_1, \{\lambda_i\}, \{\rho_i\}) = \int_{t_0}^{t_1} ch_{\check{\sigma}}(f, \{\lambda_i\}, \{\rho_i\})$.

□

A differential twisted K^1-cocycle with the twisting $\check{\sigma} = (\mathcal{G}_\sigma, \theta, \omega)$ given by a lifting bundle gerbe with connection and curving is a quadruple

$$(f, \{\lambda_i\}, \{\rho_i\}, \eta)$$

where f is a $\mathbf{PU}(\mathcal{H})$-equivariant map $\mathcal{P}_\sigma \to \mathbf{Fred}_*^{sa}$, $\{\lambda_i\}$ is a spectral cut of f with respect to an open cover $\{U_i\}$, $\{\rho_i\}$ is a partition of unity on X subordinate to $\{U_i\}$, and

$$\eta \in \Omega^{even}(X)/\mathbf{Im}(d - H),$$

where $\mathbf{Im}(d-H)$ denotes the image of the twisted de Rham operator $d-H : \Omega^{even}(X) \longrightarrow \Omega^{odd}(X)$.

We introduce an equivalence relation on the set of $\check{\sigma}$-twisted differential K^1-cocycles, generated by the following three elementary equivalence relations (See Proposition 3.3)

(1) $(f, \{\lambda_i\}, \{\rho_i\}, \eta_0)$ and $(f, \{\lambda_i\}, \{\rho'_i\}, \eta_1)$ are equivalent if $\{\lambda_i\}$ is a spectral cut of $\{f \circ \phi_i\}$ associated to local trivializing sections $\{\phi_i : U_i \to \mathcal{P}_\sigma\}$, and $\{\rho_i\}$ and $\{\rho'_i\}$ are a pair of partitions of unity on X subordinate to $\{U_i\}$, with

$$\eta_0 - \eta_1 + \eta(f, \{\lambda_i\}, \{\rho_i\}, \{\rho'_i\}) = 0 \qquad \mod (d - H).$$

(2) $(f, \{\lambda_i\}, \{\rho_i\}, \eta_0)$ and $(f, \{\mu_i\}, \{\rho_i\}, \eta_1)$ are equivalent if $\{\lambda_i\}$ and $\{\mu_i\}$ are a pair of spectral cuts for the family of operators $\{f(\phi_i(x))\}$ on an open cover $\{U_i\}$ of X, with

$$\eta_0 - \eta_1 + \eta(f, \{\lambda_i\}, \{\mu_i\}, \{\rho_i\}) = 0 \qquad \mod (d - H).$$

(3) $(f_0, \{\lambda_i\}, \eta_0)$ and $(f_1, \{\lambda_i\}, \eta_1)$ are equivalent if there exists $f : P_\sigma \times [t_0, t_1] \to \mathbf{Fred}^{sa}_*$, a $\mathbf{PU}(\mathcal{H})$-equivariant homotopy connecting $f_0 = f(\cdot, t_0)$ and $f_1 = f(\cdot, t_1)$ such that there is a family of spectral cuts $\{\lambda_i\}$ independent of $t \in [t_0, t_1]$ for $\{f(\phi_i, t)\}$ on an open cover $\{U_i \times [t_0, t_1]\}$ of $X \times [t_0, t_1]$, with

$$\eta_0 - \eta_1 + \eta(f_0, f_1, \{\lambda_i\}, \{\rho_i\}) = 0 \qquad \mod (d - H).$$

Definition 3.4. The differential twisted K-theory $\check{K}^1(X, \check{\sigma})$ with a twisting given by $\check{\sigma} = (\mathcal{G}_\sigma, \theta, \omega)$ is the space of equivalence classes of differential twisted K^1-cocycles. The differential Chern character form of a differential twisted K^1-cocycle $(f, \{\lambda_i\}, \{\rho_i\}, \eta)$ is given by

$$ch_{\check{\sigma}}(f, \{\lambda_i\}, \{\rho_i\}, \eta) = ch_{\check{\sigma}}(f, \{\lambda_i\}, \{\rho_i\}) + (d - H)\eta. \tag{3.8}$$

Theorem 3.5. *The differential Chern character form (3.8) on differential twisted K^1-cocycles defines the differential Chern character*

$$ch_{\check{\sigma}} : \check{K}^1(X, \check{\sigma}) \longrightarrow \Omega^{odd}(X, d - H),$$

whose image consists of odd degree differential forms on X, closed under $d - H$. The differential Chern character induces a well-defined twisted Chern character

$$ch_{\check{\sigma}} : K^1(X, \sigma) \longrightarrow H^{odd}(X, d - H).$$

Proof. There is a natural forgetful map $\check{K}^1(X, \check{\sigma}) \longrightarrow K^1(X, \sigma)$ which sends an equivalence class of $(f, \{\lambda_i\}, \{\rho_i\}, \eta)$ to a $\mathbf{PU}(\mathcal{H})$-equivariant homotopy class of f. Proposition 3.3 says that the differential twisted Chern character map

$$ch_{\check{\sigma}} : \check{K}^1(X, \check{\sigma}) \longrightarrow \Omega^{odd}(X, d - H)$$

induces the twisted Chern character homomorphism

$$Ch_{\check{\sigma}} : K^1(X, \sigma) \longrightarrow H^{odd}(X, H)$$

such that the following diagram commutes

$$\begin{tikzcd} \check{K}^1(X, \check{\sigma}) \arrow[r] \arrow[d, "ch_{\check{\sigma}}"] & K^1(X, \sigma) \arrow[d, "Ch_{\check{\sigma}}"] \\ \Omega^{odd}_0(X, d - H) \arrow[r] & H^{odd}(X, d - H), \end{tikzcd} \tag{3.9}$$

where $\Omega_0^{odd}(X, d-H)$ denotes the image of the differential Chern character form homomorphism $ch_{\check{\sigma}} : \check{K}^1(X, \check{\sigma}) \longrightarrow \Omega^{odd}(X, d-H)$. $\square$

It is easy to see that horizontal homomorphisms in the commutative diagram (3.9) are surjective. In order to study the kernel of the forgetful homomorphism $\check{K}^1(X, \check{\sigma}) \longrightarrow K^1(X, \sigma)$, we need to introduce the even differential twisted K-theory $\check{K}^0(X, \check{\sigma})$ and its differential Chern character. We can define a differential twisted K^0-cocycle to be a S^1-family of differential twisted K^1-cocycles which passes through a differential twisted K^1-cocycle that defines the zero element in $\check{K}^1(X, \check{\sigma})$. Then we have the following commutative diagram

$$\begin{tikzcd}
\check{K}^0(X, \check{\sigma}) \arrow[r] \arrow[d, "ch_{\check{\sigma}}"] & K^0(X, \sigma) \arrow[d, "Ch_{\check{\sigma}}"] \\
\Omega_0^{ev}(X, d-H) \arrow[r] & H^{ev}(X, d-H),
\end{tikzcd}$$

where $\Omega_0^{ev}(X, d-H)$ denotes the image of the differential Chern character homomorphism

$$ch_{\check{\sigma}} : \check{K}^0(X, \check{\sigma}) \longrightarrow \Omega^{ev}(X, d-H).$$

Theorem 3.6. *Denote by* $K^0_{\mathbb{R}/\mathbb{Z}}(X, \check{\sigma})$ *the kernel of the odd differential Chern character* $ch_{\check{\sigma}} : \check{K}^1(X, \check{\sigma}) \longrightarrow \Omega_0^{odd}(X, H)$, *then we have the following two exact sequences:*

$$0 \longrightarrow K^0_{\mathbb{R}/\mathbb{Z}}(X, \check{\sigma}) \longrightarrow \check{K}^1(X, \check{\sigma}) \xrightarrow{ch_{\check{\sigma}}} \Omega_0^{odd}(X, d-H) \longrightarrow 0, \tag{3.10}$$

$$0 \longrightarrow \frac{\Omega^{ev}(X)}{\Omega_0^{ev}(X, d-H)} \longrightarrow \check{K}^1(X, \check{\sigma}) \longrightarrow K^1(X, \sigma) \longrightarrow 0, \tag{3.11}$$

such that the following diagram commutes

(3.12)

$$\begin{tikzcd}
 & & 0 \arrow[d] & & \\
 & & K^0_{\mathbb{R}/\mathbb{Z}}(X, \check{\sigma}) \arrow[d] \arrow[dr] & & \\
0 \arrow[r] & \dfrac{\Omega^{ev}(X)}{\Omega_0^{ev}(X, d-H)} \arrow[r] \arrow[dr, "d-H"'] & \check{K}^1(X, \check{\sigma}) \arrow[r] \arrow[d, "ch_{\check{\sigma}}"] & K^1(X, \sigma) \arrow[r] \arrow[d, "Ch_{\check{\sigma}}"] & 0 \\
 & & \Omega_0^{odd}(X, d-H) \arrow[r] \arrow[d] & H^{odd}(X, d-H) & \\
 & & 0 & &
\end{tikzcd}$$

.

Proof. There is an exact sequence

$$\frac{\Omega^{ev}(X)}{\mathbf{Im}(d-H)} \longrightarrow \check{K}^1(X, \check{\sigma}) \longrightarrow K^1(X, \sigma) \longrightarrow 0,$$

where $\eta \in \dfrac{\Omega^{ev}(X)}{\mathbf{Im}(d-H)}$ is mapped to a differential twisted K^1-class given by

$$[(F, \{0\}, \{1\}, \eta)].$$

Here F is a $\mathbf{PU}(\mathcal{H})$-equivariant map from $\mathcal{P}_\sigma$ to an orbit of the adjoint action of $\mathbf{U}(\mathcal{H})$ through a fixed grading operator in $\mathcal{H}$ such that both eigenspaces are infinite dimensional. Note that $(F, \{0\}, \{1\}, \eta)$ is equivalent to $(F, \{0\}, \{1\}, 0)$ if and only if η is a differential twisted Chern character form, modulo $\mathbf{Im}(d-H)$, of an even differential twisted K^0-cocycle, that is

$$\eta \in \frac{\Omega_0^{ev}(X, d-H)}{\mathbf{Im}(d-H)}.$$

This gives rise to the exact sequence (3.11). The verification of the commutative diagram (3.12) is straightforward. □

Theorem 3.7. *Denote by* $K^1_{\mathbb{R}/\mathbb{Z}}(X, \check{\sigma})$ *the kernel of the even differential Chern character* $ch_{\check{\sigma}} : \check{K}^0(X, \check{\sigma}) \longrightarrow \Omega_0^{ev}(X, H)$, *then we have the following exact sequences:*

$$0 \longrightarrow K^1_{\mathbb{R}/\mathbb{Z}}(X, \check{\sigma}) \longrightarrow \check{K}^0(X, \check{\sigma}) \xrightarrow{ch_{\check{\sigma}}} \Omega_0^{ev}(X, H) \longrightarrow 0, \tag{3.13}$$

$$0 \longrightarrow \frac{\Omega^{odd}(X)}{\Omega_0^{odd}(X, H)} \longrightarrow \check{K}^0(X, \check{\sigma}) \longrightarrow K^0(X, \sigma) \longrightarrow 0, \tag{3.14}$$

such that the following diagram commutes

(3.15)

$$\begin{tikzcd}
 & & 0 \arrow[d] & & \\
 & & K^1_{\mathbb{R}/\mathbb{Z}}(X, \check{\sigma}) \arrow[d] \arrow[dr] & & \\
0 \arrow[r] & \dfrac{\Omega^{odd}(X)}{\Omega_0^{odd}(X, d-H)} \arrow[r] \arrow[dr, "d-H"'] & \check{K}^0(X, \check{\sigma}) \arrow[r] \arrow[d, "ch_{\check{\sigma}}"] & K^0(X, \sigma) \arrow[r] \arrow[d, "Ch_{\check{\sigma}}"] & 0 \\
 & & \Omega_0^{ev}(X, d-H) \arrow[r] \arrow[d] & H^{ev}(X, d-H) & \\
 & & 0 & . &
\end{tikzcd}$$

Remark 3.8. We point out that commutative diagrams (3.12) and (3.15) of exact sequences are joined together by two interlocking six-term exact sequences

$$\begin{tikzcd}
K^1_{\mathbb{R}/\mathbb{Z}}(X, \check{\sigma}) \arrow[r] & K^0(X, \sigma) \arrow[r, "Ch_{\check{\sigma}}"] & H^{ev}(X, d-H) \arrow[d] \\
H^{odd}(X, d-H) \arrow[u] & K^1(X, \sigma) \arrow[l, "Ch_{\check{\sigma}}"'] & K^0_{\mathbb{R}/\mathbb{Z}}(X, \check{\sigma}) \arrow[l]
\end{tikzcd}$$

and

$$\begin{tikzcd}
\dfrac{\Omega^{odd}(X)}{\Omega_0^{odd}(X, d-H)} \arrow[r, "d-H"] & \Omega_0^{ev}(X, d-H) \arrow[r] & H^{ev}(X, d-H) \arrow[d] \\
H^{odd}(X, d-H) \arrow[u] & \Omega_0^{odd}(X, d-H) \arrow[l, "Ch_{\check{\sigma}}"'] & \dfrac{\Omega^{ev}(X)}{\Omega_0^{ev}(X, d-H)} \arrow[l, "d-H"']
\end{tikzcd}$$

In [35], an analogue of the above interlocking commutative diagram in ordinary differential cohomology is called the Character Diagram.

3.2. **Classifying space of twisted K-theory.** In this subsection, we will give a universal model of twisted K-theory. We start with the even case. As a model for the classifying space of K^0 we first choose **Fred**, the space of all bounded Fredholm operators in an infinite dimensional separable complex Hilbert space $\mathcal{H}$. Consider the twisted product

$$\mathfrak{G} = \mathbf{PU}(\mathcal{H}) \ltimes GL(\infty),$$

where the group $\mathbf{PU}(\mathcal{H})$ acts on the group $GL(\infty)$, of operators that are invertible and 1+ finite rank, by conjugation. Thus the product in $\mathfrak{G}$ is given by

$$(g, f) \cdot (g', f') = (gg', f(gf'g^{-1})).$$

A principal $\mathfrak{G}$-bundle over X, locally trivial with respect to an open cover $\{U_i\}$, is defined by transition functions $(g_{ij}, f_{ij}) : U_i \cap U_i \longrightarrow \mathfrak{G}$ satisfying the usual cocycle condition which encodes both the cocycle relation for the transition functions $\{g_{ij}\}$ of the $\mathbf{PU}(\mathcal{H})$ bundle over X and the twisted cocycle relation for $\{f_{ij}\}$

$$f_{ij}(\hat{g}_{ij} f_{jk} \hat{g}_{ij}^{-1}) = f_{ik}, \tag{3.16}$$

which is independent of the choice of the lifting $\hat{g}_{ij}$ of g_{ij} to $\mathbf{U}(\mathcal{H})$.

Next we construct a universal $\mathfrak{G}$ bundle E over the classifying space $B\mathfrak{G} = E/\mathfrak{G}$. The total space is just the Cartesian product $E = \mathcal{P} \times \mathcal{Q}$, where $\mathcal{P}$ is the total space of a universal $\mathbf{PU}(\mathcal{H})$ bundle over the base $K(\mathbb{Z}, 3)$ and $\mathcal{Q}$ is the total space of a universal $GL(\infty)$ bundle over the base $\mathbf{Fred}^{(0)}$, the index zero component of the classifying space of K^0. The total space of $\mathcal{Q}$ is the set of pairs (q, w) with q a parametrix of a index zero Fredholm operator w. This space is contractible. This follows from the observation that the pairs can be parametrized by (q, t) with $w = (1+t)q^{-1}$ and t is an arbitrary finite rank operator, q an arbitrary invertible operator. The right action of $GL(\infty)$ is given by $(q, w) \cdot a = (qa, w)$ and is free.

The right action of $(g, a) \in \mathfrak{G}$ on E is defined by

$$(p, (q, w)) \cdot (g, a) = (pg, (g^{-1}qag, g^{-1}wg))$$

where w is a Fredholm operator with parametrix q. It is easy to see that the action is free, because $\mathbf{PU}(\mathcal{H})$ acts freely on the first component and the second component involves right multiplication of the invertible operator q by the group element $a \in GL(\infty)$.

Now we can establish a model for the classifying space for even twisted K-theory.

Theorem 3.9. *Given a principal* $\mathbf{PU}(\mathcal{H})$*-bundle* $\mathcal{P}_\sigma$ *over* X *with* $\sigma : X \to K(\mathbb{Z}, 3)$ *representing a non-torsion class in* $H^3(X, \mathbb{Z})$*, then the even twisted K-theory* $K^0(X, \sigma)$ *can be identified with the set of homotopy classes of maps* $X \to B\mathfrak{G}$ *covering the map* σ*.*

Proof. Let $f : \mathcal{P}_\sigma \to \mathbf{Fred}$ be a $\mathbf{PU}(\mathcal{H})$-equivariant family of Fredholm operators. We can select an open cover $\{U_i\}$ of X such that on each U_i there is a local section $\phi_i : U_i \to \mathcal{P}_\sigma$ and for each i the index zero Fredholm operators $f(\phi_i(x))$, $x \in U_i$ have a gap in the spectrum at some $\lambda_i \neq 0$. Then over U_i we have a finite rank vector bundle E_i defined by the spectral projection $f(\phi_i(x))^2 < \lambda_i^2$. As usual the transition functions are $g_{ji} : U_{ji} \to \mathbf{PU}(\mathcal{H})$ with $\phi_i(x) = \phi_j(x) g_{ji}(x)$ and the lifts $\hat{g}_{ij}$ of g_{ij} to $\mathbf{U}(\mathcal{H})$ satisfy $\hat{g}_{ij}\hat{g}_{jk}\hat{g}_{ki} = \sigma_{ijk} Id$ on triple intersections with σ_{ijk} taking values in $U(1)$.

We may assume that E_i is a trivial vector bundle over U_i of rank n_i by passing to a finer cover if necessary. Choosing a trivialization of E_i gives a $\mathbb{Z}_2$ graded parametrix $q_i(x)$ (an inverse up to finite rank operators) for each $f(\phi_i(x))$, $x \in U_i$. The operator $q_i(x)^{-1}$ is defined as the direct sum of the restriction of $f(\phi_i(x))$ to the orthogonal complement of E_i in $\mathcal{H}$ and an isomorphism between the vector bundles E_i^+ and E_i^-. Clearly then $f(\phi_i(x)) q_i(x) = 1$ modulo rank n_i operators.

For $x \in U_{ij}$ we have a pair of parametrices $q_i(x)$ and $q_j(x)$ for $f(\phi_i(x))$ and $f(\phi_j(x))$ respectively. These are related by an invertible operator $f_{ij}(x)$ of type 1+ finite rank and

$$\hat{g}_{ij} q_j(x) \hat{g}_{ij}^{-1} = q_i(x) f_{ij}(x).$$

The conjugation on the left hand side by $\hat{g}_{ij}$ comes from the equivariance relation

$$f(\phi_j(x)) = f(\phi_i(x) g_{ij}(x)) = \hat{g}_{ij}(x)^{-1} f(\phi_i(x)) \hat{g}_{ij}(x).$$

The system $\{f_{ij}\}$ does not quite satisfy the Čech cocycle relation because of the different local sections $\phi_i : U_i \to \mathcal{P}_\sigma$ involved. Instead, we have on U_{ijk}

$$\hat{g}_{jk} q_k \hat{g}_{jk}^{-1} = q_j f_{jk} = (\hat{g}_{ij}^{-1} q_i f_{ij} \hat{g}_{ij}) f_{jk} = \hat{g}_{jk} (\hat{g}_{ik}^{-1} q_i f_{ik} \hat{g}_{ik}) \hat{g}_{jk}^{-1}.$$

Using the relation $\hat{g}_{jk}\hat{g}_{ik}^{-1} = \sigma_{ijk}\hat{g}_{ij}^{-1}$, we get

$$\hat{g}_{jk} (\hat{g}_{ik}^{-1} q_i f_{ik} \hat{g}_{ik}) \hat{g}_{jk}^{-1} = \hat{g}_{ij}^{-1} q_i f_{ik} \hat{g}_{ij}$$

multiplying the last equation from right by $\hat{g}_{ij}^{-1}$ and from the left by $q_i^{-1}\hat{g}_{ij}$ one gets the twisted cocycle relation

$$f_{ij} (\hat{g}_{ij} f_{jk} \hat{g}_{ij}^{-1}) = f_{ik},$$

which is independent of the choice of the lifting $\hat{g}_{ij}$. We may think of (3.16) as defining an untwisted cocycle relation for $\{(g_{ij}, f_{ij})\}$ in the twisted product $\mathfrak{G} = \mathbf{PU}(\mathcal{H}) \ltimes GL(\infty)$, In summary, this cocycle $\{(g_{ij}, f_{ij})\}$ defines a principal $\mathfrak{G}$ bundle over X whose classifying map is a continuous map $X \to B\mathfrak{G}$. A homotopic $\mathbf{PU}(\mathcal{H})$-equivariant family of Fredholm operators gives rise to a homotopic classifying map $X \to B\mathfrak{G}$. □

Remark 3.10. The base space $B\mathfrak{G}$ is a fiber bundle over $K(\mathbb{Z}, 3)$. The projection is defined by

$$\xi((p, (q, w))\mathfrak{G}) = \pi(p),$$

where $\pi : \mathcal{P} \to K(\mathbb{Z}, 3)$ is the projection. The fiber $\xi^{-1}(z)$ at $z \in K(\mathbb{Z}, 3)$ is isomorphic (but not canonically so) to the space **Fred** of Fredholm operators; to set up the isomorphism one needs a choice of element p in the fiber $\pi^{-1}(z)$. Thus a section of the bundle $B\mathfrak{G}$ over $K(\mathbb{Z}, 3)$ is the same thing as a $\mathbf{PU}(\mathcal{H})$ equivariant map γ from $\mathcal{P}$ to **Fred**. We use the notation $K^0(K(\mathbb{Z}, 3), \mathcal{P})$ for the twisted K-group of $K(\mathbb{Z}, 3)$ with $\mathcal{P}$ being thought of as the twist. Then γ is defining an element of $K^0(K(\mathbb{Z}, 3), \mathcal{P})$. This latter group is not known (and of course the same is true for the odd twisted K-group of $K(\mathbb{Z}, 3)$). The case of ordinary K-theory on the Eilenberg-MacLane space $K(\mathbb{Z}, 3)$ is already complicated, but it is known that it is given by the integral cohomology of the space $K(\mathbb{Q}, 3)$, (Cf. [1]).

The twisted Chern character on $K^0(X, \sigma)$ can also be constructed by choosing a connection ∇ on a principal $\mathfrak{G}$ bundle over X associated to the cocycle $\{(g_{ij}, f_{ij}\}$. Locally, on a trivializing open cover $\{U_i\}$ of X, we can lift the connection to a connection taking values in the Lie algebra $\hat{\mathbf{g}}$ of the central extension $U(H) \times GL(\infty)$ of $\mathfrak{G}$. Denote by $\hat{F}_\nabla$ the curvature of this connection. On the overlaps U_{ij} the curvature satisfies a twisted relation

$$\hat{F}_{\nabla,j} = Ad_{(g_{ij}, f_{ij})^{-1}} \hat{F}_{\nabla,i} + g_{ij}^* c,$$

where c is the curvature of the canonical connection θ on the principal $U(1)$-bundle $\mathbf{U}(\mathcal{H}) \to \mathbf{PU}(\mathcal{H})$.

Since the Lie algebra $\mathbf{u}(\infty) \oplus \mathbb{C}$ is an ideal in the Lie algebra of $U(\mathcal{H}) \times GL(\infty)$, the projection $F'_{\nabla,i}$ of the curvature $\hat{F}_{\nabla,i}$ onto this subalgebra transforms in the same way as $\hat{F}$ under change of local trivialization. It follows that for a $\mathbf{PU}(\mathcal{H})$-equivariant map $f : \mathcal{P}_\sigma \to \mathbf{Fred}$, we can define a twisted Chern character form of f as

$$ch_{\check{\sigma}}(f, \nabla) = e^{B_i} \mathrm{tr}\, e^{F'_{\nabla,i}/2\pi i},$$

over U_i. Here the trace is well-defined on $\mathbf{gl}(\infty)$ and on the center $\mathbb{C}$ it is defined as the coefficient of the unit operator. Note that $ch_{\check{\sigma}}(f, \nabla)$ is globally defined and $(d - H)$-closed

$$(d - H) ch_{\check{\sigma}}(f, \nabla) = 0,$$

and depends on the differential twisting $\check{\sigma} = (\mathcal{G}_\sigma, \theta, \omega)$, and a choice of a connection ∇ on a principal $\mathfrak{G}$ bundle over X.

We now give a short description of the classifying space in odd twisted K-theory. First let $\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-$ be a decomposition into isomorphic subspaces. Operators on $\mathcal{H}$ can be written as two by two matrices of operators with respect to this decomposition. The group $\mathbf{U}_{res} = \mathbf{U}_{res}(\mathcal{H})$ is the group of unitary operators in $\mathcal{H}$ with Hilbert-Schmidt off-diagonal blocks (for more on this definition see [33]). In the non-twisted case an element of $K^1(X)$ is given by a homotopy class of maps $X \to U(\infty)$. Since $U(\infty)$ is the base space of a universal $\mathbf{U}_{res}$ bundle, this is the same thing as giving an equivalence class of $\mathbf{U}_{res}(\mathcal{H})$ bundles over X.

Remark 3.11. The universal $\mathbf{U}_{res}$ bundle Q can be constructed as follows. By Bott periodicity, $\mathbf{U}_{res}$ is homotopy equivalent [33] to the group $\Omega U(\infty)$ of smooth based loops in $U(\infty)$. The universal $\Omega U(\infty)$ bundle Q over $U(\infty)$ is simply the space of smooth paths

$f : [0,1] \to U(\infty)$ with $f(0) = 1$ and $f^{-1}df$ periodic. The right action of $\Omega U(\infty)$ on Q is just the pointwise multiplication of paths, see Appendix 2 in [13].

A $\mathbf{U}_{res}$ bundle over X can be given in terms of local transition functions $\phi_{ij} : U_{ij} \to \mathbf{U}_{res}$. Next we need to twist this construction by the transition functions g_{ij} of the $\mathbf{PU}(\mathcal{H})$ bundle over X. In the odd case for any $g \in \mathbf{PU}(\mathcal{H})$ we had an automorphism Aut_g of $U(\infty)$ given by a conjugation by any unitary operator covering $g \in \mathbf{PU}(\mathcal{H})$. In the case of $\mathbf{U}_{res}$ we have to be a little more careful since conjugation by a unitary operator is not in general an automorphism of $\mathbf{U}_{res}$. However, we recall that $\mathbf{U}_{res}$ is a classifying space for even K-theory and is homotopy equivalent to the space of bounded Fredholm operators in any separable infinite dimensional Hilbert space. Actually, the homotopy equivalence is given explicitly as the map $g \mapsto a$, where

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

in a decomposition $\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-$, [33], [41]. The block a is always a Fredholm operator and one can construct a homotopy inverse to this map. This gives directly the right way to define automorphisms of $\mathbf{U}_{res}$ by elements of $\mathbf{PU}(\mathcal{H}_+)$, compatible with automorphisms of the space of Fredholm operators: For any $g \in \mathbf{PU}(\mathcal{H}_+)$ one selects $\hat{g} \in \mathbf{U}(\mathcal{H}_+)$ and considers this as a unitary operator in $\mathcal{H}$ by

$$g \mapsto \begin{pmatrix} \hat{g} & 0 \\ 0 & \hat{g} \end{pmatrix}.$$

Thus, in the same way as in the even case, we can think of an element in the twisted K-theory group $K^1(X, \sigma)$ defined by transition functions $h_{ij} : U_{ij} \to \mathfrak{H}$, where $\mathfrak{H}$ is the group

$$\mathfrak{H} = \mathbf{PU}(\mathcal{H}_+) \times_{\mathbf{PU}(\mathcal{H}_+)} \mathbf{U}_{res}$$

with multiplication $(g, h)(g', h') = (gg', h(gh'g^{-1})$.

More precisely, using the concrete realization of the universal $\mathbf{U}_{res}$ bundle in Remark 3.11, we can choose local sections $q_i : U_i \to Q$ by choosing for each $x \in U_i$ a path $q_i(x)$ in $U(\infty)$ joining the neutral element to the end point $f(\phi_i(x))$, where we have taken f as the equivariant function on $\mathcal{P}_\sigma$ taking values in the model $U(\infty)$ for the classifying space $\mathbf{Fred}_*^{sa}$ of odd K-theory. On the intersection U_{ij} we then have

$$\hat{g}_{ij}(x) q_j(x) \hat{g}_{ij}^{-1}(x) = q_i(x) \cdot f_{ij}(x),$$

where $f_{ij}(x)$ is an element of the loop group $\Omega U(\infty)$, which by the Remark gives an element in $\mathbf{U}_{res}$. From this transformation rule we obtain the group multiplication law above for the transition functions $h_{ij} = (g_{ij}, f_{ij})$ taking values in the group $\mathfrak{H}$.

As in the even case, one can check that the classifying space $B\mathfrak{H}$ is a fiber bundle over the base $K(\mathbb{Z}, 3)$ with $\mathbf{Fred}_*^{sa}$ (or equivalently, $U(\infty)$) as the fiber. It is written as $B\mathfrak{H} = \mathcal{P} \times_{\mathbf{PU}(\mathcal{H}_+)} \mathbf{Fred}_*^{sa}$, where now $\mathcal{P}$ is the universal $\mathbf{PU}(\mathcal{H}_+)$ bundle over $K(\mathbb{Z}, 3)$.

Theorem 3.12. *Given a principal* $\mathbf{PU}(\mathcal{H}_+)$*-bundle* $\mathcal{P}_\sigma$ *over* X *defined by* $\sigma : X \to K(\mathbb{Z}, 3)$*, the odd twisted K-group* $K^1(X, \sigma)$ *is the set of homotopy classes of maps* $X \to B\mathfrak{H}$ *covering the map* σ*.*

Example 3.13. Let E be a real Euclidean vector bundle of rank $2n$ over X. Then there is a lifting bundle gerbe $\mathcal{G}_{W_3(E)}$ over X, called the Spin bundle gerbe, associated to the frame bundle $SO(E)$ and the flat $U(1)$-bundle over $SO(2n)$:

$$U(1) \to Spin^c(2n) = Spin(2n) \times_{\mathbb{Z}_2} U(1) \longrightarrow SO(2n).$$

Choose a local trivialization of E over an open cover $\{U_i\}$ of X. Then the transition functions

$$g_{ij} : U_i \cap U_j \longrightarrow SO(2n).$$

define an element in $H^1(X, \underline{SO(2n)})$ whose image under the Bockstein exact sequence

$$H^1(X, \underline{Spin(2n)}) \to H^1(X, \underline{SO(2n)}) \to H^2(X, \mathbb{Z}_2)$$

is the second Stieffel-Whitney class $w_2(E)$ of E. The corresponding $\mathbf{PU}(\mathcal{H})$-bundle is defined by the following twisting

$$W_3(E) := W_3 \circ \nu_E : X \longrightarrow \mathbf{BSO}(2n) \longrightarrow K(\mathbb{Z}, 3),$$

where ν_E is the classifying map of E and W_3 is the classifying map of the principal $U(1)$-bundle $\mathbf{BSpin}^c(2n) \to \mathbf{BSO}(2n)$.

Note that the Spin bundle gerbe can be equipped with a flat connection and trivial curving. Denote also by $w_2(E)$ the corresponding bundle gerbe with a flat connection and a trivial curving. With respect to a good cover $\{U_i\}$ of X the differential twisting $w_2(E)$ defines a Deligne cocycle $(\sigma_{ijk}, 0, 0)$ with trivial local B-fields, here $\sigma_{ijk} = \hat{g}_{ij}\hat{g}_{jk}\hat{g}_{ki}$ where $\hat{g}_{ij} : U_{ij} \to Spin^c(2n)$ is a lift of g_{ij}.

A differential twisted K-class in $\check{K}^0(X, w_2(E))$ can be represented by a Clifford bundle, denoted $\mathcal{E}$, equipped with a Clifford connection, where E is also equipped with a $SO(2n)$-connection. Locally, over each U_i we let $\mathcal{E}|_{U_i} \cong S_i \otimes \mathcal{E}_i$ where S_i is the local fundamental spinor bundle associated to $E|_{U_i}$ with the standard Clifford action of $\mathbf{Cliff}(E|_{U_i})$ obtained from the fundamental representation of $Spin(2n)$. Then $\mathcal{E}_i$ is a complex vector bundle over U_i with a connection ∇_i such that

$$ch(\mathcal{E}_i, \nabla_i) = ch(\mathcal{E}_j, \nabla_j).$$

Hence, our construction of a differential Chern character implies that the twisted Chern character

$$Ch_{w_2(E)} : K^0(X, W_3(E)) \longrightarrow H^{ev}(X)$$

is given by sending $[\mathcal{E}]$ to $\{[ch(\mathcal{E}_i, \nabla_i)] = ch(\mathcal{E}_i)\}$. If E is equipped with a $Spin^c$ structure whose determinant line bundle is L, there is a canonical isomorphism

$$K^0(X) \longrightarrow K^0(X, W_3(E)),$$

given by $[V] \mapsto [V \otimes S_E]$ where S_E is the associated spinor bundle of E. Then we have

$$Ch_{w_2(E)}([V \otimes S_E]) = e^{\frac{c_1(L)}{2}} ch([V]),$$

where $ch([V])$ is the ordinary Chern character of $[V] \in K^0(X)$.

If X is an even dimensional Riemannian manifold, and TX is equipped with the Levi-Civita connection, we can identify $K^0(X, W_3(E))$ with the Grothendieck group of Clifford modules, denoted $K^0(X, \mathbf{Cliff}(TX))$, of the bundle of Clifford algebras $\mathbf{Cliff}(TX)$. Then

$$Ch_{w_2(X)}([\mathcal{E}]) = ch(\mathcal{E}/S) \tag{3.17}$$

where $ch(\mathcal{E}/S)$ is the relative Chern character of the Clifford module $\mathcal{E}$ constructed in Section 4.1 of [6]. In particular, if X is equipped with a $Spin^c$ structure (Cf. [26]), let its canonical class be c_1, and its complex spinor bundle associated to the fundamental representation of $Spin^c(2n)$ be $S(X)$. Then any Clifford module $\mathcal{E}$ can be written as $V \otimes S(X)$, whose twisted Chern character is given by

$$Ch_{w_2(X)}(\mathcal{E}) = e^{\frac{c_1}{2}} ch(V),$$

where $ch(V)$ is the ordinary Chern character. Note that the vector bundle V in the decomposition $\mathcal{E} = V \otimes S(X)$ depends on the choice of $Spin^c$ structure, but $e^{\frac{c_1}{2}} ch(V)$ depends only on $\mathcal{E}$.

4. RIEMANN-ROCH THEOREM IN TWISTED K-THEORY

A smooth map $f : X \to Y$ is called K-oriented if $TX \oplus f^*TY$ is equipped with a Spin^c structure which is determined by a choice of $c_1 \in H^2(X, \mathbb{Z})$ such that

$$w_2(X) - f^* w_2(Y) = c_1 \mod 2.$$

The push-forward map $f_!^{c_1} : K^*(X) \to K^{*-d(f)}(Y)$, also called the Gysin homomorphism in K-theory, is well-defined, here $d(f) = (\dim Y - \dim X) \mod 2$. The Riemann-Roch theorem (Cf. [2]) is given by the following formula

$$Ch\big(f_!^{c_1}(a)\big)\hat{A}(Y) = f_*^H\big(Ch(a) e^{\frac{c_1}{2}} \hat{A}(X)\big), \tag{4.1}$$

where $a \in K^*(X)$, and f_*^H is the Gysin homomorphism in ordinary cohomology theory, Ch is the Chern character for ordinary complex K-theory, $\hat{A}(X)$ and $\hat{A}(Y)$ are the A-hat classes (which can be expressed in terms of the Pontrjagin classes) of X and Y respectively.

Let $W_3(f)$ represent the image of $f^*(w_2(Y)) - w_2(X)$ under the Bockstein homomorphism $H^2(X, \mathbb{Z}_2) \longrightarrow H^3(X, \mathbb{Z})$. In [16], we showed that there is a natural push-forward map (2.4)

$$f_!^K : K^i\big(X, f^*\sigma + W_3(f)\big) \longrightarrow K^{i+d(f)}(Y, \sigma),$$

associated to any differentiable map $f : X \to Y$ and any $\sigma : Y \to K(\mathbb{Z}, 3)$. The push-forward map is constructed by choosing a closed embedding $\iota : X \to Y \times \mathbb{R}$ such that $f = \pi \circ \iota$ where π is the projection. The push-forward is functorial in the sense that

$$f_!^K = \pi_!^K \circ \iota_!^K,$$

where the push-forward map $\pi_!^K$ is given by the Bott periodicity of twisted K-theory. So we may assume that $f : X \to Y$ is a closed embedding and X and Y are equipped with compatible Riemannian metrics.

In this Section we will establish the Riemann-Roch theorem in twisted K-theory for any closed embedding $f : X \to Y$ with a twisting $\sigma : Y \to K(\mathbb{Z}, 3)$ (the classifying map for $\mathcal{P}_\sigma$). We do this in two stages, represented by Theorems 4.1 and 4.4. Theorem 4.1 is a special case (that we need in this paper) which we will prove by a method that may be easily modified to give the general case (Theorem 4.3). Recall the notation $\check{\sigma} = (\mathcal{G}_\sigma, \theta, \omega)$ for a differential twisting and H for the normalized curvature of $\check{\sigma}$. We use $w_2(f)$ to denote the Spin bundle gerbe with a trivial connection and a trivial curving on X associated to the normal bundle N_f of the embedding f (Cf. Example 3.13).

We first assume that $f : X \to Y$ is K-oriented, that is, the normal bundle $N_f \cong f^*TY/TX$ of f is equipped with a *Spin*^{c} structure. Under this assumption, we have the following Riemann-Roch theorem in twisted K-theory for a K-oriented embedding $f : X \to Y$ and $[f^*\sigma] = 0$. For simplicity, we assume that $\dim Y - \dim X = 0 \mod 2$, otherwise, we can replace Y by $Y \times \mathbb{R}$.

Theorem 4.1. *Let $f : X \to Y$ be a closed embedding whose normal bundle is equipped with a Spin^c structure. Assume that $\dim Y - \dim X = 2n$ and $f^*[\sigma] = 0$. Let $a \in K^0(X, f^*\sigma)$, and $c_1(f)$ be the canonical class of the Spin^c structure on N_f. Let f_*^H be the Gysin homomorphism in twisted cohomology theory, and $Ch_{\check{\sigma}}$ and $Ch_{f^*\check{\sigma}}$ be the twisted Chern characters on $K^0(Y, \sigma)$ and $K^0(X, f^*\sigma)$ respectively. Then*

$$Ch_{\check{\sigma}}\big(f_!^K(a)\big)\hat{A}(Y) = f_*^H\big(Ch_{f^*\check{\sigma}}(a)e^{\frac{-c_1(f)}{2}}\hat{A}(X)\big).$$

Proof. Choose a local trivialization of $\mathcal{P}_\sigma$ which defines a degree two Deligne cocycle $(\sigma_{ijk}, A_{ij}, B_i)$ on Y associated to $\check{\sigma}$ with respect to an open cover $\{U_i\}$ and transition functions $g_{ij} : U_i \cap U_j \longrightarrow \mathbf{PU}(\mathcal{H})$. The conditions $f^*[\sigma] = [W_3(f)] = 0$ imply that there is a homotopy commutative diagram (Cf. [37])

$$\begin{tikzcd} X \arrow[r, "\nu_f"] \arrow[d, "f"'] & \mathbf{BSO}(2n) \arrow[d, "W_3"] \arrow[dl, Rightarrow, "\eta"'] \\ Y \arrow[r, "\sigma"'] & K(\mathbb{Z}, 3). \end{tikzcd} \tag{4.2}$$

Here ν_f is the classifying map of the normal bundle of f, W_3 is the classifying map of the principal $BU(1)$-bundle $\mathbf{BSpin}^c(2n) \to \mathbf{BSO}(2n)$ and η is a homotopy between $W_3 \circ \nu_f$ and $\sigma \circ f$.

The homotopy commutative diagram (4.2) and the *Spin*^{c} structure on N_f define a trivialization of $f^*\mathcal{P}_\sigma$ given by a section $\psi : X \to f^*\mathcal{P}_\sigma$. With respect to the induced open cover from $\{U_i\}$ and the trivialization of $f^*\mathcal{P}_\sigma$, the transition functions $\{g_{ij}\}$ can be written as

$$g_{ij} = h_i h_j^{-1}$$

for $h_i : \tilde{U}_i = X \cap U_i \to \mathbf{PU}(\mathcal{H})$. Then the pull-back differential twist $f^*\check{\sigma}$ defines a degree 2 Deligne cocycle

$$(1, A_j - A_i, B_i)$$

such that $B_i - dA_i$ is a globally defined 2-form on X, denoted by B_X, and $dB_X = f^*H$ on X. Note that such a Deligne cocycle can be extended to a tubular neighborhood of X in Y with the 2-form denoted by B_f.

Given a twisted K-class in $K^0(X, f^*\sigma)$, represented by a locally defined map $\psi_i : \tilde{U}_i \to$ $\mathbf{Fred}(\mathcal{H})$, we have

$$h_i^{-1}\psi_i h_i = h_j^{-1}\psi_j h_j,$$

which implies that $\{h_i^{-1}\psi_i h_i\}$ defines a continuous map $\psi : X \to \mathbf{Fred}(\mathcal{H})$. Our construction of the differential twisted Chern character shows that

$$ch_{f^*\check{\sigma}}(\{\psi_i\}) = \exp(B_X) ch(\psi)$$

where ch is the ordinary Chern character form of ψ. Hence we have established the existence of the following commutative diagram relating the ordinary Chern character on $K^0(X)$ and the twisted Chern character on $K^0(X, f^*\sigma)$ respectively

(4.3)
$$\begin{tikzcd} K^0(X) \arrow[r, "\cong"] \arrow[d, "Ch"] & K^0(X,\ f^*\sigma) \arrow[d, "Ch_{f^*\check{\sigma}}"] \\ H^{ev}(X, d) \arrow[r, "\exp(B_X)", "\cong"'] & H^{ev}(X, d - f^*H). \end{tikzcd}$$

Note that the above arguments also work for N_f which is identified with the tubular neighborhood of X in Y. Denote by $\iota : N_f \to Y$ the open embedding, so we have a similar commutative diagram relating the ordinary Chern character on $K^0(N_f)$ and the twisted Chern character on $K^0(N_f, \iota^*\sigma)$ respectively

(4.4)
$$\begin{tikzcd} K^0(N_f) \arrow[r, "\cong"] \arrow[d, "Ch"] & K^0(N_f, \iota^*\sigma) \arrow[d, "Ch_{\iota^*\check{\sigma}}"] \\ H^{ev}(N_f, d) \arrow[r, "\exp(B_f)", "\cong"'] & H^{ev}(N_f, d - \iota^*H). \end{tikzcd}$$

For the open embedding $\iota : N_f \to Y$, the twisted Chern characters commute with the push-forward map $\iota_!^K$, so the following diagram commutes

(4.5)
$$\begin{tikzcd} K^0(N_f, \iota^*\sigma) \arrow[r, "\iota_!^K"] \arrow[d, "Ch_{\iota^*\check{\sigma}}"] & K^0(Y, \sigma) \arrow[d, "Ch_{\check{\sigma}}"] \\ H^{ev}(N_f, d - \iota^*H) \arrow[r, "\iota_!^H"'] & H^{ev}(Y, d - H). \end{tikzcd}$$

Introduce the zero section $o : X \to N_f$, so that $f = \iota \circ o$, and the Riemann Roch theorem for o gives what we call the 'Chern character defect diagram'

(4.6)
$$\begin{tikzcd} K^0(X) \arrow[r, "o_!^K"] \arrow[d, "Ch"] & K^0(N_f) \arrow[d, "Ch"] \\ H^{ev}(X, d) \arrow[r, "\cong", "o_!"'] & H^{ev}(N_f, d), \end{tikzcd}$$

which is not commutative. Here the Gysin homomorphism $o_!$ is the Thom isomorphism. The non-commutativity of (4.6) is described by the following formula

$$Ch\big(o_!^K(a)\big)\hat{A}(N_f) = o_!^H\big(Ch(a)e^{\frac{-c_1(f)}{2}}\hat{A}(X)\big), \tag{4.7}$$

for any $a \in K^0(X)$ where the homomorphism $o_!^H$ on twisted cohomology theory is

$$o_!^H = \exp(B_f) \circ o_! \circ \exp(-B_X) : H^{ev}(X, d - f^*H) \longrightarrow H^{ev}(N_f, d - \iota^*H).$$

The Gysin homomorphism $f_!^H$ in twisted cohomology theory is given by $\iota_!^H \circ o_!^H$.

The commutative diagrams (4.3), (4.4), (4.5) and the formula (4.7) imply that

$$Ch_{\check{\sigma}}\big(f_!^K(a)\big)\hat{A}(Y) = f_*^H\big(Ch_{f^*\check{\sigma}}(a)e^{\frac{-c_1(f)}{2}}\hat{A}(X)\big),$$

for any $a \in K^0(X, f^*\sigma)$. Here we have applied the identity $\hat{A}(N_f) = \iota^*\hat{A}(Y)$. □

Remark 4.2. In ordinary K-theory, for a closed embedding $f : X \to Y$ whose normal bundle is equipped with a *Spin*^{c} structure $c_1(f)$, then f is $-c_1(f)$-oriented as in [2], and Theorem 4.1 agrees with the Atiyah-Hirzebruch's Riemann-Roch theorem.

Remark 4.3. Under the assumptions of Theorem 4.1, for a closed Riemannian embedding $f : X \to Y$ whose normal bundle is equipped with a Hermitian structure, we have an integral version of the Riemann-Roch Theorem 4.1. To describe this, let E be an Hermitian vector bundle with Hermitian connection ∇^E over X. Let ∇^{TX} and ∇^{TY} denote the Levi-Civita connections on TX and TY respectively, and ∇_f be an Hermitian connection on N_f. Then

$$\int_Y ch_{\check{\sigma}}\big(f_!^K(E, \nabla^E)\big)\hat{A}(Y, \nabla^{TY}) = \int_X ch(E, \nabla^E)e^{\frac{-c_1(N_f, \nabla_f)}{2}}\hat{A}(X, \nabla^{TX}), \tag{4.8}$$

The equality (4.8) follows from our construction of the twisted Chern character form and the differential version of the diagrams (4.3), (4.4), (4.5) and the formula (4.7).

The proof of Theorem 4.1 and arguments in Example 3.13 can be adapted to give the following general Riemann-Roch theorem in twisted K-theory. We leave the proof to dedicated readers.

Theorem 4.4. *Let* $f : X \to Y$ *be a closed embedding. Assume that* $\dim Y - \dim X = 2n$. *Then*

$$Ch_{\check{\sigma}}\big(f_!^K(a)\big)\hat{A}(Y) = f_*^H\big(Ch_{f^*\check{\sigma}+w_2(f)}(a)\hat{A}(X)\big),$$

for any $a \in K^0(X, f^*\sigma + W_3(f))$. *Equivalently, the following diagram commutes*

$$\begin{tikzcd} K^0(X, f^*\sigma + W_3(f)) \arrow[r, "f_!^K"] \arrow[d, "Ch_{f^*\check{\sigma}+w_2(f)}(-)\hat{A}(X)"'] & K^0(Y, \sigma) \arrow[d, "Ch_{\check{\sigma}}(-)\hat{A}(Y)"] \\ H^{ev}(X, f^*H) \arrow[r, "f_*^H"'] & H^{ev}(Y, H). \end{tikzcd} \tag{4.9}$$

In particular, if $\sigma = W_3(Y)$, *then the Riemann-Roch formula is given by*

$$Ch_{w_2(Y)}\big(f_!^K(a)\big)\hat{A}(Y) = f_*^H\big(Ch_{w_2(X)}(a)\hat{A}(X)\big).$$

for any $a \in K^0(X, W_3(X))$.

5. Applications

In this Section, we apply our differential twisted K-theory and Riemann-Roch theorem to investigate twisted K-theory of simply connected simple Lie groups. From the result of Douglas ([17]) and Braun ([10]), we know that the twisted K-theory of a simply connected simple Lie group of rank n is an exterior algebra on $(n-1)$ generators tensor a cyclic group. The order of the cyclic group can be described in terms of the dimensions of irreducible representations of G. So the twisted Chern characters are all zero for the twisted K-theory of a simply connected simple Lie group. We will apply twisted Chern character forms to study these torsion elements in twisted K-theory.

Let G be a simply connected compact simple Lie group of rank n and T be a maximal torus of G. The sets of simple roots and fundamental weights of G with respect to T are denoted by

$$\{\alpha_1, \alpha_2 \cdots, \alpha_n\}, \qquad \{\lambda_1, \lambda_2 \cdots, \lambda_n\}$$

respectively. Let $\mathbf{k}$ denote the twisting $G \to K(\mathbb{Z}, 3)$ corresponding to the integer k in $H^3(G, \mathbb{Z}) \cong \mathbb{Z}$ with the corresponding $\mathbf{PU}(\mathcal{H})$ bundle $\mathcal{P}_{\mathbf{k}}$.

In [15], a canonical trivialization of the associated bundle gerbe $\mathcal{G}_{\mathbf{k}}$ over certain conjugacy classes in G is constructed. Those conjugacy classes are diffeomorphic to G/T, called symmetric D-branes in [8], [18], [19], [23] and labeled by dominant weights of level k. For each simple root α_i, the corresponding subgroup $SU(2)_{\alpha_i}$ intersects with these D-branes at $S^2_{\alpha_i}$, which splits $SU(2)_{\alpha_i}$ into two 3-dimensional balls D_i^+ and D_i^-. Denote by $C_i = C_i^+ \cup C_i^-$ the CW complex obtained by attaching these two 3-dimensional cells D_i^+ and D_i^- to G/T.

Let $\iota_i : C_i \to G$ be the inclusion map. Then the restriction $\iota_i^* \mathbf{k} = \mathbf{k} \circ \iota_i : C_i \to K(\mathbb{Z}, 3)$ defines the integer k in $H^3(C_i, \mathbb{Z}) \cong \mathbb{Z}$. Note that $C_i^+ \cap C_i^- = G/T$, and denote by $\iota_i^\pm : G/T \to C_i^\pm$ and $\iota : G/T \to G$ the inclusion maps. The twisting k is trivial when restricted to $C_i^\pm$, and the two trivializations on G/T differ by a line bundle $L_{k\lambda_i}$ whose first Chern class is given by $k\lambda_i \in H^2(G/T)$. We have the following Mayer-Vietoris sequence

(5.1)
$$\begin{tikzcd}
K^1(C_i^+) \oplus K^1(C_i^-) \arrow[r] & 0 \arrow[r] & K^0(C_i, \mathbf{k}) \arrow[d] \\
K^1(C_i, \mathbf{k}) \arrow[u] & K^0(G/T) \arrow[l, "\delta"'] & K^0(C_i^+) \oplus K^0(C_i^-) \arrow[l]
\end{tikzcd}$$

where the map $K^0(C_i^+) \oplus K^0(C_i^-) \to K^0(G/T)$ sends $([E_1], [E_2])$ to $[E_1] - [L_{k\lambda_i} \otimes E_2]$. This implies that the push-forward map

$$\iota_!^K : K^0(G/T) \longrightarrow K^{\dim G}(G, \mathbf{k})$$

satisfies $\iota_!^K([E]) = \iota_!^K([L_{k\lambda_i} \otimes E])$ for any vector bundle E over G/T.

Choose a gerbe connection and a curving on $\mathcal{P}_{\mathbf{k}}$ to get a differential twisting $\check{\mathbf{k}}$ whose normalized curvature is H. As the twisted Chern characters are zero on $K^*(G, \mathbf{k})$, we see that

$$K^*(G, \mathbf{k}) \cong K^{*+1}(G, \mathbf{k}, \mathbb{R}/\mathbb{Z}),$$

which is a subgroup of differential twisted K-theory $\check{K}(G, \check{\mathbf{k}})$ by Theorems 3.6 and 3.7, under a homomorphism sending $[f]$ to an element $[f, \eta] \in \check{K}(G, \check{\mathbf{k}})$ such that $ch_{\check{\mathbf{k}}}(f) = -(d-H)\eta$. Therefore, the differential twisted Chern character of $\iota^K_!(E)$, for a complex vector bundle E over G/T, has the form $(d-H)\eta(E)$ for some differential form $\eta(E)$. Then we have

$$\int_G (d-H)\eta(E) = \int_G (d-H)\eta(L_{k\lambda_i} \otimes E) \mod c_G(k)$$

where $c_G(k)$ is the cyclic order of $K^{\dim G}(G, \mathbf{k})$.

As G/T is a complex manifold, the normal bundle of G/T in G has a canonical $Spin^c$ structure, we have the following integral version of our Riemann-Roch Theorem 4.1 (Cf. Remark 4.3)

$$\int_G (d-H)\eta(E) = \int_{G/T} Ch(E)\mathbf{Td}(G/T) \tag{5.2}$$

where $\mathbf{Td}(G/T)$ is the Todd class of G/T. Hence, we obtain

$$\int_{G/T} Ch(E)\mathbf{Td}(G/T) = \int_{G/T} Ch(L_{k\lambda_i} \otimes E)\mathbf{Td}(G/T) \mod c_G(k),$$

for $i = 1, \cdots, n$. In particular, let E be a trivial line bundle, then we have the following relations

$$\int_{G/T} Ch(L_{k\lambda_i})\mathbf{Td}(G/T) = 1 \mod c_G(k), \tag{5.3}$$

for $i = 1, \cdots, n$, where the left hand side is the dimension of the irreducible representation of G with highest weight $k\lambda_i$. We can repeat the above arguments by attaching 3-dimensional cells along various $S^2_{\alpha_i}$ simultaneously and we obtain the following general relations

$$\int_{G/T} ch(L_{k\lambda_{i_1}+k\lambda_{i_2}+\cdots+k\lambda_{i_m}})\mathbf{Td}(G/T) = 1 \mod c_G(k), \tag{5.4}$$

for any subset $\{i_1, i_2, \cdots, i_m\}$ of $\{1, 2, \cdots, n\}$. Denote by $V(k\lambda_{i_1} + k\lambda_{i_2} + \cdots + k\lambda_{i_m})$ the irreducible representation of G with highest weight $k\lambda_{i_1} + k\lambda_{i_2} + \cdots + k\lambda_{i_m}$, then we have the following identities relating the cyclic order of $K^{\dim G}(G, \mathbf{k})$ to the highest weight irreducible representations of G

$$\dim V(k\lambda_{i_1} + k\lambda_{i_2} + \cdots + k\lambda_{i_m}) = 1 \mod c_G(k), \tag{5.5}$$

for any subset $\{i_1, i_2, \cdots, i_m\}$ of $\{1, 2, \cdots, n\}$.

5.1. **The $SU(2)$ case.** In this subsection $X = SU(2) = S^3$. It is sufficient to consider an open cover U_0, U_1 consisting of slightly extended hemispheres with an intersection homotopic to S^2. In this case it was shown in [34] that $K^1(SU(2), \mathbf{k}) \cong \mathbb{Z}_k$, and the cyclic order agrees with the relation given by (5.5).

In the twisted cocycle of vector bundles we have now only a single element

$$(E_{01}, \nabla_{01}) \longrightarrow U_{01}$$

with no condition on the vector bundle. The only topological information are the rank n_{01} of the vector bundle (which could be negative for formal differences of vector bundles) and the total degree on U_{01}, which corresponds to an integer m times the Chern class of the basic complex line bundle on S^2, represented by the first Chern form

$$\omega_{01}^{[2]} = c_1(E_{01}, \nabla_{01}).$$

The Chern character form of (E_{01}, ∇_{01}) modulo the Chern character of

$$(E_1, \nabla_1) - (E_0, \nabla_0) \otimes L_{01},$$

where L_{01} is a complex line bundle on the intersection of degree k, is then equal to $\mathbb{Z}_k$ since both E_0 and E_1 is trivial and they are characterized by the ranks n_0, n_1. This is actually just the standard argument using the Mayer-Vietoris sequence for two open sets.

Choosing a partition of unity ρ_0, ρ_1 subordinate to the open cover U_i and forms B_i with

$$dB_i = H = kH_0, \qquad B_0 - B_1 = c_{01}$$

on the overlap $U_{01} = U_0 \cap U_1$, where H_0 is the normalized volume form on $SU(2) = S^3$, we obtain the the $(d-H)$-exact form on $SU(2)$ representing the twisted Chern differential character form Θ,

$$\begin{aligned}\Theta_0 &= (d-H)[\rho_1 n_{10} + \rho_1(\omega_{10}^{[2]} + B_0 n_{10})] \\ &= n_{10} d\rho_1 + d\rho_1 \wedge \omega_{10}^{[2]} + n_{10} d\rho_1 \wedge B_0.\end{aligned}$$

on U_0, and

$$\begin{aligned}\Theta_1 &= (d-H)[\rho_0 n_{01} + \rho_0(\omega_{01}^{[2]} + B_1 n_{01})] \\ &= n_{01} d\rho_0 + d\rho_0 \wedge \omega_{01}^{[2]} + n_{01} d\rho_0 \wedge B_1.\end{aligned}$$

on U_1. Note that $n_{01} = -n_{10}$, $d\rho_0 + d\rho_1 = 0$ and $\omega_{10}^{[2]} + B_0 n_{10} = -(\omega_{01}^{[2]} + B_1 n_{01})$ imply that $\Theta_0 = \Theta_1$ on U_{01}. As $H^{odd}(X, d-H) = 0$, we know that

$$\Theta = (d-H)(\eta_{[0]} + \eta_{[2]})$$

for a globally defined even form $(\eta_{[0]} + \eta_{[2]})$, called a twisted eta potential. We will show that these twisted eta potentials are localized to certain conjugacy classes such that

$$\int_{SU(2)} \eta_{[0]} H_0 = \frac{m}{k} \mod 1$$

using the Riemann-Roch theorem in twisted K-theory (Theorem 4.1).

A Fredholm operator realization of the twisted K-theory classes in $K^1(SU(2), \mathbf{k})$ is obtained from the family of hamiltonians in a supersymmetric Wess-Zumino-Witten model, [29]. Actually in this case all the classes can be realized as equivariant twisted K-theory classes, equivariant under the conjugation action of $SU(2)$ on itself. We recall some basic properties of the Fredholm family from [30], [21].

We have self-adjoint Fredholm operators Q_A in a fixed Hilbert space $\mathcal{H}$ parametrized by $SU(2)$ connections A (termed 'vector potentials' in the physics literature) on the unit circle S^1. These transform equivariantly under gauge transformations,

$$\hat{g}^{-1} Q_A \hat{g} = Q_{A^g}$$

where g is an element of the loop group LG, $\hat{g}$ is a projective representation of the loop group in $\mathcal{H}$, i.e., a representation of the standard central extension of level k, and $A^g = g^{-1}Ag + g^{-1}dg$ is the gauge transformed vector potential.

Let $p : \mathcal{A} \to G$ be the canonical projection from the space of vector potentials on the circle to the group of holonomies around the circle; the fiber of this projection is the group of based loops $\Omega G \subset LG$. The spectrum of Q_A depends only on the projection $p(A) \in G$, by the equivariance property.

For certain A the operators Q_A have kernels. Those A for which this is true are such that $p(A)$ is a conjugacy class $\mathcal{C}_j$, the so called 'D-brane', in G. The physics terminology is to say that the zero modes of the family are localised in $\mathcal{C}_j$.

The conjugacy class is diffeomorphic to the sphere S^2 given by

$$\mathcal{C}_j = \{ghg^{-1} | g \in G\}$$

where $h = e^{i\pi \frac{2j+1}{k}\sigma_3}$ with $\sigma_3 = diag(1, -1)$ and $2j = 0, 1, 2, \dots k-2$. The zero modes form a complex line bundle L_j over $\mathcal{C}_j$ with Chern class represented by $2j+1$ times the basic 2-form on $S^2 \cong \mathcal{C}_j$.

For the later example it is worth noticing here that under the map $p : \mathcal{A} \to G$, the conjugacy classes correspond to coadjoint orbits in the Lie algebra of the central extension of the loop algebra; the space of vector potentials on the circle is the (non-centrally extended) loop algebra, the coadjoint action corresponds to the gauge action on $\mathcal{A}$.

If we consider a non-zero eigenvalue λ of Q_A we can still conclude from the continuity of the family Q_A as a function of A that for sufficiently small values of $|\lambda|$ the eigenvectors corresponding to this eigenvalue are localized at a 2-sphere close to the 2-sphere $\mathcal{C}_j$ defined by the zero modes and the eigenvectors corresponding to λ form a complex line bundle of winding number $2j+1$. Thus if we fix a small real number $0 < \epsilon$ the spectral subspace $-\epsilon < Q_A < \epsilon$ is a complex line in a tubular neighborhood $\tilde{\mathcal{C}_j}$ of $\mathcal{C}_j$.

Setting

$$U_\pm = \{g \in G | \pm\epsilon \notin \mathbf{Spec}(Q_A), A \in p^{-1}(g)\},$$

the intersection $U_- \cap U_+$ consists of $\tilde{\mathcal{C}_j}$ and two contractible components $D_\pm$ (upper and lower hemispheres in S^3). In this case we have only one spectral vector bundle E_{-+} which is the extension of L_j to the tubular neighborhood $\tilde{\mathcal{C}_j}$ and a trivial line bundle on the remaining components $D_\pm$ of $U_- \cap U_+$. The constraint $0 < 2j+1 < k$ is compatible with the known result $K^1(SU(2), \mathbf{k}) = \mathbb{Z}/k\mathbb{Z}$. The construction gives a realization for all elements except the neutral element in $K^1(SU(2), \mathbf{k})$. The neutral element can then be obtained for example as the sum of classes corresponding to $2j+1 = 1$ and $2j+1 = k-1$.

Write the normalized curvature H of $\check{\mathbf{k}}$ as kH_0, where H_0 represents the basic 3-form on $SU(2)$. As the twisted cohomology $H^{odd}(SU(2), d - kH_0) = 0$, instead we study the differential twisted Chern character form by applying the Riemann-Roch theory (Theorem 4.1).

By (5.2), we have

$$\begin{aligned}\int_{SU(2)}(d-H)(\eta_{[0]}+\eta_{[2]}) &= \int_{\mathcal{C}_j} c_1(L_j,\nabla_j) \mod k \\ &= (2j+1) \mod k.\end{aligned}$$

Thus we have, as $H = kH_0$,

$$\int_{SU(2)} \eta_{[0]} H_0 = -\frac{2j+1}{k} \mod 1.$$

Hence, the twisted eta potential distinguishes the twisted K-classes in $K^1(SU(2), \mathbf{k})$.

5.2. **The $SU(3)$ case.** As the final example we study the odd K-group $K^1(SU(3), \mathbf{k})$. We need a representation of the twisted affine Lie algebra $A_2^{(2)}$. Here the twist refers to an outer automorphism τ of $\mathbf{su(3)}$ with $\tau^2 = 1$. Then this algebra is defined as a central extension of the subalgebra $L_\tau G$ consisting of smooth maps $g : [0, \pi] \to \mathbf{su(3)}$ such that $g(\pi) = \tau(g(0))$. In the case of $SU(3)$ we can take τ to be given by complex conjugation of matrices. This automorphism of the Lie algebra integrates to an automorphism of the group $SU(3)$, again given by complex conjugation.

The loop algebra of $SO(3)$ is clearly a subalgebra of $L_\tau G$ and the central extension of the former on level k is obtained as a restriction of the central extension of level k of the latter.

The gauge conjugation action sending $Q_A \to \hat{g}^{-1} Q_A \hat{g} = Q_{A^g}$ is now given by the coadjoint right action $A \mapsto A^g$ where the 'vector potential' A is an element in the dual $L_\tau \mathbf{su(3)}^*$. The coadjoint orbits have been analyzed in [38] and are shown to be equal to the so called *twisted conjugacy classes* in $SU(3)$. Recall that in this terminology a twisted conjugacy class $C(h)$ corresponding to $h \in SU(3)$ is defined as

$$C(h) = \{gh\tau(g)^{-1} | g \in SU(3)\}.$$

The general formula for the (twisted) conjugacy class corresponding to the zero modes of the operators Q_A is $h = \exp[-2\pi(\lambda^\vee + \rho^\vee)/k]$ where ρ is half the sum of positive roots of the Lie algebra of constant gauge transformations (which in this case is $\mathbf{so(3)}$) and λ is the highest weight of an irreducible $\mathbf{so(3)}$ representation, [21]. Here $\lambda^\vee \in \mathbf{h}$ is the dual of a weight $\lambda \in \mathbf{h}^*$, the duality is determined by the Killing form. Note also that the level $k \geq \kappa$, where κ is the dual Coxeter number (in the case of $SU(n)$ this number is equal to n). The reason for this is that in the Wess-Zumino-Witten model construction $k = k' + \kappa$ where k' is the level of an arbitrary irreducible loop group representation and the shift κ comes from a loop group representation constructed from the Clifford algebra of the loop group.

As shown in [36] the twisted conjugacy classes C_x defined by $h(x) = \exp 2x\rho^\vee$ define a foliation of $SU(3)$ when $0 \leq x \leq \pi/4$. Concretely, as a 3×3 matrix,

$$h(x) = \begin{pmatrix} \cos(2x) & \sin(2x) & 0 \\ -\sin(2x) & \cos(2x) & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

When $x = 0$ the twisted conjugacy class can be identified as the 5-dimensional space $M_5 = SU(3)/SO(3)$, when $x = \pi/4$ it is $S^5 = SU(3)/SU(2)$ whereas for $0 < x < \pi/4$ we obtain the 'seven brane' $M_7 = SU(3)/SO(2)$.

The class defined by the zero modes of Q_A is the generic orbit M_7 corresponding to the parameter value $x = (2j+1)/k$, where $2j = 0, 1, 2 \ldots k-3$, the shift by 3 coming from the dual Coxeter number of $SU(3)$. In the case of the twisted loop algebra $A_2^{(2)}$ we have an additional constraint: The spin j is integer when $k-3$ is even and j is a half-integer when $k-3$ is odd.

The degree of the zero mode line bundle is computed as in the $SU(2)$ case. The zero modes are localized in the finite-dimensional vacuum subspace of the Hilbert space $H_{j,k}$ which carries a representation $j \otimes 1/2$ of $\mathbf{so(3)}$. (In general, the zero mode bundle has rank equal to the multiplicity of the representation ρ inside of the spin representation of the Clifford algebra of the group of constant loops; here however the Clifford algebra representation is two dimensional and at the same time the irreducible fundamental representation of the Lie algebra $\mathbf{so(3)}$.) When $x = (2j+1)/k$ the complex line is spanned by the highest weight vector $v_{j+1/2}$ of $\mathbf{so(3)}$ weight $j + 1/2$, [21]. The zero mode bundle is then the associated line bundle $L = SU(3) \times_{2j+1} \mathbb{C}$, defined by the principal bundle $SO(2) \to SU(3) \to M_7$ through the one dimensional representation of $SO(2)$ with character $2j+1$. In particular, for even k the spin j is a half-integer and thus the degree of the zero mode bundle is even. This is in accordance with the known result $K^1(SU(3), \mathbf{k}) = \mathbb{Z}/(k/2)\mathbb{Z}$ for even k. Twisted K^1 is a rank one module over the untwisted $K^0(SU(3))$, tensoring with any of the elements in $K^1(SU(3), \mathbf{k})$ gives only even elements in $\mathbb{Z}_k$.

In the case when k is odd, the degree of the zero mode bundle is also odd and the tensor product operation $K^0(SU(3)) \times K^1(SU(3), \mathbf{k}) \to K^1(SU(3), \mathbf{k})$, gives both the even and odd elements in $\mathbb{Z}_k$.

The eta forms detect the twisted K-theory classes in a similar way as in the case of $SU(2)$. In the present setting we have nonzero eta forms in even degrees up to form degree six. However, they do not contain independent information since for any fixed level k the only parameter is the twisting of the line bundle L which is given by the integer $2j+1$. To determine this integer from the differential data one proceeds as in the case of $SU(2)$ above. Consider the subgroup $SU(2) \subset SU(3)$ given by the the matrices with $+1$ on the diagonal in the lower right corner. The intersection of $SU(2) = S^3$ with the orbit M_7 is a union of two spheres S^2_a and S^2_b. The reason for this is that the points $h(x)$ and $h(-x)$ are conjugate to each other by the twisted action of $g = diag(1, -1, -1)$ which lies outside of $SU(2)$; one can see easily that $h(\pm x)$ are not conjugate by elements of $SU(2)$ and their union is $M_7 \cap SU(2)$. So the differential twisted Chern character form is given by

$$(d - H)(\eta_{[0]} + \cdots + \eta_{[6]}).$$

Picking up the component of form degree 3 we get

$$\begin{aligned}\int_{SU(2)} \eta_{[0]} H &= \textstyle\int_{S^2} c_1(L_j, \nabla_j) \mod c_G(k) \\ &= 2(2j+1) \mod c_G(k),\end{aligned}$$

the factor 2 coming from the integration of c_1 over the two distinct 2-spheres. Thus again, we get

$$\int_{SU(2)} \eta_{[0]} H_0 = -\frac{2(2j+1)}{k} \mod c_G(k)/k.$$

References

[1] D.W. Anderson and L. Hodgkin: The K-theory of Eilenberg-MacLane complexes, Topology, 7, 317-329, 1968.

[2] M. Atiyah and F. Hirzebruch, *Riemann-Roch theorems for differentiable manifolds.* Bull. Amer. Math. Soc. 65 276–281, 1959.

[3] M. Atiyah and G. Segal, *Twisted K-theory.* Ukr. Math. Bull. Vol. 1, 291–334, 2004.

[4] M. Atiyah and G. Segal, *Twisted K-theory and cohomology.* Preprint, math.KT/0510674.

[5] Atiyah, M. F. and Singer, I. M. *Index theory for skew-adjoint Fredholm operators*, Inst. Hautes tudes Sci. Publ. Math. No. 37 5–26, 1969.

[6] N. Berline, E. Getzler and M. Vergne, *Heat kernels and Dirac operators,* Grundlehren Math. Wiss., vol. 298, Springer-Verlag, New York, 1992.

[7] P. Bouwknegt, A. Carey, V. Mathai, M. Murray and D. Stevenson, *Twisted K-theory and K-theory of bundle gerbes.* Comm. Math. Phys., 228, 17–45, 2002.

[8] P. Bouwknegt, P. Dawson and D. Ridout, *D-branes on group manifolds and fusion rings.* JHEP 0212. 065, 2002.

[9] P. Bouwknegt, J. Evslin and V. Mathai, *T-Duality: Topology Change from H-flux.* Commun.Math.Phys. 249, 383-415, 2002.

[10] V. Braun, *Twisted K-Theory of Lie Groups*, arXiv:hep-th/0305178. JHEP 0403, 029, 2004

[11] J-L, Brylinski, *Loop spaces, characteristic classes and geometric quantization,* Progress in Mathematics, 107. Birkhauser Boston, 1993.

[12] U. Bunke and T. Schick, *Smooth K-theory*, preprint, arXiv:0707.0046.

[13] A. L. Carey and J.Mickelsson, *A gerbe obstruction to quantization of fermions on odd-dimensional manifolds with boundary,* Lett. Math. Phys. Vol. 51, 145–160, 2000.

[14] A. L. Carey and J. Phillips, *Unbounded Fredholm modules and spectral flow.* Canad. J. Math. 50, 673–718, 1998.

[15] A. L. Carey and B.L. Wang, *Fusion of Symmetric D-branes and Verlinde ring.* Comm. in Math. Phys., 277, 577-625, 2008. math.ph/0505040.

[16] A. L. Carey and B.L. Wang, *Thom isomorphism and Push-forward map in twisted K-theory.* Journal of K-theory. math.KT/0507414

[17] C. Douglas, *On the Twisted K-Homology of Simple Lie Groups*, arXiv:math/0402082.

[18] G. Felder, J. Frohlich, J. Fuchs and C. Schweigert, *The geometry of WZW branes.* J. Geom. Phys. 34, 162-190, 2000.

[19] S. Fredenhagen and V. Schomerus, *Branes on Group Manifolds, Gluon Condensates, and twisted K-theory.* JHEP 0104, 007, 2001.

[20] D. Freed, *Dirac charge quantization and generalized differential cohomology.* Surveys in differential geometry, 129–194, Surv. Differ. Geom., VII, Int. Press, Somerville, MA, 2000.

[21] D. S. Freed, Michael J. Hopkins and C. Teleman, *Twisted K-theory and loop group representations* arXiv:math/0312155

[22] D. Freed, E. Witten, *Anomalies in String Theory with D-Branes*, hep-th/9907189.

[23] K. Gawedzki and N. Reis, *WZW branes and gerbes.* Rev. Math. Phys. 14, 1281-1334, 2002.

[24] M.J. Hopkins and I.M. Singer, *Quadratic functions in geometry, topology,and M-theory*, preprint, math.AT/0211216

[25] N. Kuiper, *The homotopy type of the unitary group of Hilbert space.* Topology, 3, 19-30, 1965.

[26] B. Lawson and M-L. Michelsohn, *Spin Geometry*, Princeton University Press, 1989.

[27] J. Lott, $\mathbb{R}/\mathbb{Z}$ *index theory.* Comm. Anal. Geom. 2, 279–311,1994.

[28] V. Mathai and D. Stevenson, *Chern character in twisted K-theory: equivariant and holomorphic cases*, Commun.Math.Phys. 228, 17-49, 2002.

[29] J. Mickelsson, *Gerbes, (twisted) K-theory, and the supersymmetric WZW model*, hep-th/0206139. Infinite dimensional groups and manifolds, 93–107, IRMA Lect. Math. Theor. Phys., 5, de Gruyter, Berlin, 2004.

[30] J. Mickelsson and J.P. Pellonpaa, *Families index theorem in supersymmetric WZW model and twisted K-theory: The SU(2) case.*, preprint, hep-th/0509064.

[31] M. K. Murray *Bundle gerbes,* J. London Math. Soc. (2) **54**, 403–416, 1996.

[32] M. K. Murray and D. Stevenson, *Bundle gerbes: stable isomorphism and local theory.* J. London Math. Soc. (2) 62 (2000), no. 3, 925–937.

[33] A. Pressley and G. Segal, *Loop Groups*, Clarendon Press, Oxford (1986)

[34] J. Rosenberg, *Continuous trace* C^**-algebras from the bundle theoretic point of view*, J. Aust. Math. Soc. **A47**, 368-384 1989.

[35] J. Simons and D.Sullivan, *Axiomatic characterization of ordinary differential cohomology.* Jour. of Topology 1, 45–56 2008.

[36] S. Stanciu, *An illustrated guide to D-branes in SU(3)*, arXiv:hep-th/0112221

[37] B.L. Wang, *Geometric cycles, Index theory and twisted K-homology.* Preprint, arXiv:0710.1625.

[38] R. Wendt, *Twisted conjugacy classes, coadjoint orbits of loop groups and D-branes in the WZW-model*, arXiv:math/0303118

[39] E. Witten, *D-branes and K-theory* J. High Energy Phys. 12, Paper 19, 1998.

[40] E. Witten, *Overview of K-theory applied to strings*, Int. J. Mod. Phys. **A16**, 693, 2001, hep-th/0007175.

[41] T. Wurzbacher, *An elementary proof of the homotopy equivalence between the restricted general linear group and the space of Fredholm operators* In: Analysis, geometry and topology of elliptic operators, 411–426, World Sci. Publ., Hackensack, NJ (2006)

(Alan L. Carey) Mathematical Sciences Institute, Australian National University, Canberra ACT 0200, Australia

E-mail address: `acarey@maths.anu.edu.au`

(Jouko Mickelsson) Department of Mathematics and Statistics, University of Helsinki, Finland, and Department of Theoretical Physics, KTH, Stockholm, Sweden

E-mail address: `jouko.mickelsson@helsinki.fi`

(Bai-Ling Wang) Department of Mathematics, Mathematical Sciences Institute, Australian National University, Canberra, ACT 0200, Australia.

E-mail address: `wangb@maths.anu.edu.au`